U.S. DEPARTMENT OF COMMERCE
National Technical Information Service
PB80-122260

Freshwater Algae of Rae Lakes Basin Kings Canyon National Park (California)

(U.S.) Environmental Monitoring and Support Lab, Las Vegas, NV

Jul 79

United States
Environmental Protection
Agency

Environmental Monitoring
and Support Laboratory
P.O. Box 15027
Las Vegas NV 89114

EPA-600/3-79-080
July 1979

PB80-122260

Research and Development

EPA

Freshwater Algae of Rae Lakes Basin, Kings Canyon National Park

REPRODUCED BY
NATIONAL TECHNICAL
INFORMATION SERVICE
U.S. DEPARTMENT OF COMMERCE
SPRINGFIELD, VA. 22161

RESEARCH REPORTING SERIES

Research reports of the Office of Research and Development, U.S. Environmental Protection Agency, have been grouped into nine series. These nine broad categories were established to facilitate further development and application of environmental technology. Elimination of traditional grouping was consciously planned to foster technology transfer and a maximim interface in related fields. The nine sereies are:

1. Environmental Health Effects Research
2. Environmental Protection Technology
3. Ecological Research
4. Environmental Monitoring
5. Socioeconomic Environmental Studies
6. Scientific and Technical Assessment Reports (STAR)
7. Interagency Energy—Environment Research and Development
8. "Special" Reports
9. Miscellaneous Reports

This report has been assigned to the ECOLOGICAL RESEARCH series. This series describes research on the effects of pollution on humans,plant and animal species, and materials. Problems are assessed for their long-and short-term influences. Investigations include formations, transport, and pathway studies to determine the fate of pollutants and their effects. This work provided the technical basis for setting standards to minimize undesirable changes in living organisms in the aquatic, terrestrial, and atmospheric environments.

This document is available to the public through the National Technical Information Service, Springfield, Virginia 22161

TECHNICAL REPORT DATA
(Please read Instructions on the reverse before completing)

1. REPORT NO.	2.	3. RECIPIENT'S ACCESSION NO.
EPA-600/3-79-080		PB 80 122260

4. TITLE AND SUBTITLE	5. REPORT DATE
FRESHWATER ALGAE OF RAE LAKES BASIN, KINGS CANYON NATIONAL PARK	July 1979
	6. PERFORMING ORGANIZATION CODE
7. AUTHOR(S)	8. PERFORMING ORGANIZATION REPORT NO.
William D. Taylor	
9. PERFORMING ORGANIZATION NAME AND ADDRESS	10. PROGRAM ELEMENT NO.
Environmental Monitoring and Support Laboratory Office of Research and Development U.S. Environmental Protection Agency Las Vegas, NV 89114	1BD884
	11. CONTRACT/GRANT NO.
12. SPONSORING AGENCY NAME AND ADDRESS	13. TYPE OF REPORT AND PERIOD COVERED
U.S. Environmental Protection Agency-Las Vegas, NV Office of Research and Development Environmental Monitoring and Support Laboratory Las Vegas, NV 89114	8-16-72 to 8-25-72
	14. SPONSORING AGENCY CODE EPA/600/07

15. SUPPLEMENTARY NOTES

16. ABSTRACT

This report illustrates and characterizes algae (exclusive of diatoms) found in Kings Canyon National Park, California and describes their distribution among the Rae Lakes within. It is the first taxonomic study of the freshwater algae for the southern Sierra Nevada and the most comprehensive for the range. It serves as a reference manual for the identification of algae in alpine and subalpine regions and establishes baseline data for future investigations. More than half (113) of the 210 forms encountered were desmids (Chlorophyta). While 120 forms were thought to be new records for California, one variety was thought to be new to science. A table illustrating the distribution of taxa within the lakes and ponds is included and discussed.

17. KEY WORDS AND DOCUMENT ANALYSIS

a. DESCRIPTORS	b. IDENTIFIERS/OPEN ENDED TERMS	c. COSATI Field/Group
Algae Freshwater Taxonomy	alpine, subalpine habitat Sierra Nevada range algal survey	06 C 08 H

18. DISTRIBUTION STATEMENT	19. SECURITY CLASS *(This Report)*	21. NO. OF PAGES
RELEASE TO PUBLIC	UNCLASSIFIED	119
	20. SECURITY CLASS *(This page)*	22. PRICE
	UNCLASSIFIED	PC: A06 / MF A01

EPA Form 2220-1 (Rev. 4-77) PREVIOUS EDITION IS OBSOLETE

* U.S. GOVERNMENT PRINTING OFFICE 1979- 683-091/2210

EPA-600/3-79-080
July 1979

FRESHWATER ALGAE OF RAE LAKES BASIN,

KINGS CANYON NATIONAL PARK

by

William D. Taylor

Water and Land Quality Branch
Monitoring Operations Division
Environmental Monitoring and Support Laboratory
Las Vegas, Nevada 89114

U.S. ENVIRONMENTAL PROTECTION AGENCY
OFFICE OF RESEARCH AND DEVELOPMENT
ENVIRONMENTAL MONITORING AND SUPPORT LABORATORY
LAS VEGAS, NEVADA 89114

DISCLAIMER

This report has been reviewed by the Environmental Monitoring and Support Laboratory-Las Vegas, U.S. Environmental Protection Agency, and approved for publication. Mention of trade names or commercial products does not constitute endorsement or recommendation for use.

FOREWORD

Protection of the environment requires effective regulatory actions that are based on sound technical and scientific information. This information must include the quantitative description and linking of pollutant sources, transport mechanisms, interactions, and resulting effects on man and his environment. Because of the complexities involved, assessment of specific pollutants in the environment requires a total systems approach that transcends the media of air, water, and land. The Environmental Monitoring and Support Laboratory-Las Vegas contributes to the formation and enhancement of a sound monitoring data base for exposure assessment through programs designed to:

- develop and optimize systems and strategies for monitoring pollutants and their impact on the environment
- demonstrate new monitoring systems and technologies by applying them to fulfill special monitoring needs of the Agency's operating programs

This report illustrates and characterizes algae found in the Kings Canyon National Park and describes their distribution among the Rae Lakes within. It is the first algal taxonomic study for the southern Sierra Nevada and the most comprehensive for the range. It serves as a reference manual for the identification of algae in alpine and subalpine regions and establishes baseline data for future investigations. This report was written for use by Federal, State, and local governmental agencies concerned with the application of biological data to water quality analysis and monitoring. Private industry and individuals similarly involved with the biological aspects of water quality will find the document useful. For further information contact the Water and Land Quality Branch, Monitoring Operations Division.

George B. Morgan
Director
Environmental Monitoring and Support Laboratory
Las Vegas

ABSTRACT

This report illustrates and characterizes algae (exclusive of diatoms) found in Kings Canyon National Park, California and describes their distribution among the Rae Lakes within. It is the first taxonomic study of the freshwater algae for the southern Sierra Nevada and the most comprehensive for the range. It serves as a reference manual for the identification of algae in alpine and subalpine regions and establishes baseline data for future investigations. More than half (113) of the 210 forms encountered were desmids (Chlorophyta). While 120 forms were thought to be new records for California, one variety was thought to be new to science. A table illustrating the distribution of taxa within the lakes and ponds is included and discussed.

CONTENTS

Foreword iii
Abstract iv
Figures and Tables vi
Acknowledgments vii

Introduction 1
Description of the Rae Lakes Area 2
    Geography 2
    Lakes and Ponds 2
Materials and Methods 6
    Collection 6
    Preparation and Examination 7
Results and Discussion 8

Literature Cited 20
Appendices 24

    A. Species, Variety and Form Descriptions 24
    B. Species Illustrations 65

Genus and Species Index 106

FIGURES

Number		Page
1.	Rae Lakes, Kings Canyon National Park, California.	3

TABLES

Number		Page
1.	Descriptive Summary of the Rae Lakes and Ponds During August 16-25, 1972, and Samples Collected	5
2.	Summary of the Taxa Encountered	9
3.	Distribution of Algal Species by Lake and Pond	11

ACKNOWLEDGMENTS

Many of the data presented in this report were a portion of a thesis submitted to Humboldt State University in partial fulfillment of the requirements for the degree of Master of Arts, July 1974. I am greatly indebted to Dr. William C. Vinyard, my major professor, for direction and guidance on this study and for the collection of the samples upon which this report is based. Dr. Vinyard collected the samples as part of an intensive investigation he conducted into the possible eutrophication of the Rae Lakes through overuse of the region by hikers. I would also like to thank Dr. G. W. Prescott, University of Montana Biological Station, for his interest and response to inquiries during the completion of this project.

INTRODUCTION

Eutrophication, i.e., the process by which water bodies become nutrient-rich, is of great significance to water quality managers. One of the primary symptoms of eutrophication is the excessive growth of algae. This growth is used to monitor the trophic state or condition of lakes. In order to use algae as a biological water quality monitoring tool, it is necessary to be able to identify them and to have information on their distribution and habitat requirements.

The purpose of this study was to list and illustrate the freshwater algae (exclusive of diatoms) of the Rae Lakes Basin, a subalpine area of the Sierra Nevada. I am aware of only three reports dealing with the freshwater algal flora of the Sierra Nevada: Wolle 1887, Vinyard 1951 (unpublished master's thesis), and Thomasson 1962. The general lack of taxonomic publications covering this large geographic area emphasizes the need for further information.

DESCRIPTION OF THE RAE LAKES AREA

GEOGRAPHY

Kings Canyon National Park is located on the broad western slope of the Sierra Nevada, occupying 1833 square kilometers. The mountains in this region range in elevation from about 427 meters (m) in the western foothills to 4,418 m at Mount Whitney on the eastern crest. The region was glaciated several times in the Pleistocene epoch, resulting in an area of high relief and exposed granite which is representative of much of the Sierra Nevada.

The Rae Lakes (lat. 36°49' N., long. 118°24' W.) are in the eastern section of the park near the main crest at an altitude of 3,231 m (Figure 1). They are accessible only on foot or by horseback, requiring about two days travel time. The Dragon pluton, the principal rock type forming the lake basins, is a quartz monzonite-granodiorite that is poor in quartz (Moore 1963).

LAKES AND PONDS

The Rae Lake system consists of six interconnected alpine and subalpine lakes along a 4 kilometer (km) stretch of valley floor and two hanging-cirque lakes 150 to 180 meters above the general elevation of the other lakes (Figure 1). The basin runs north-south, with steep sides rising over 762 m to the surrounding peaks. Snowmelt is the primary source of water for these lakes. The water flows from south to north throughout the system and forms headwaters for the Kings River watershed.

Dr. W. C. Vinyard, of Humboldt State University, who took samples in the area August 16-25, 1972, noted that the two southernmost lakes were the largest. Lake 1, a cirque lake, was bordered by rock and barren of any conspicuous vegetation. Lake 2 had a maximum depth of 16 m. Coleman (1925) reported a maximum depth of 16.5 m in the Rae Lakes but failed to specify the lake to which the measurement applied. Aquatic vegetation in Lake 2 was limited to a well-developed marsh in the shallow northeast bay. Lake 3 was shallow; Lake 4 was essentially a Carex marsh; and Lakes 5 and 6 were larger, apparently with more depth than Lakes 3 and 4. Time constraints prevented sampling visits to the two hanging-cirque lakes, Dragon Lake and the unnamed lake located on the western wall of the canyon.

Two small ponds were included in the study. Pond A was between and to the east of Lakes 2 and 3. It was about 15 centimeters (cm) deep. Pond B was located between Lakes 5 and 6 and its depth was unknown.



Figure 1. Rae Lakes, Kings Canyon National Park, California.

A descriptive summary of the study lakes and ponds is given in Table 1. At the time of collection the lakes and ponds were slightly acid, ranging in pH from 5.8 to 6.6; water temperatures ranged from 14.4 to 18.9 degrees Celsius.

The benthic vegetation was dominated by Elodea nuttalii forming dense beds at depths greater than 1 m. Elodea nuttalii was present in Lakes 2 and 6 and extended into the outlet streams. Nitella sp., a macroscopic chlorophyte, grew in dense beds fringing E. nuttalii in water less than one m deep. Neither plant was found in Lake 1. A sparse population of Ranunculus aquatilia was observed at the margin and in the outlet stream of Lake 6 and within the E. nuttalii beds in the stream between Lakes 5 and 6. Isoetes bolanderi was occasionally found in Lakes 2 and 3; Sparganium sp. and Callitriche sp. covered half of pond A.

The observed emergent vegetation was restricted to Carex sp. dominating a small marsh in a cove at the northeast corner of Lake 2 and nearly surrounding Lake 4. Otherwise, the plant was scattered.

The terrestrial vegetation included several species of Salix growing along streams and near lake shores. Pinus contorta and P. balfouriana were found intermixed, scattered, or in small clumps throughout the basin.

TABLE 1. DESCRIPTIVE SUMMARY OF THE RAE LAKES AND PONDS DURING AUGUST 16 - 25, 1972, AND SAMPLES COLLECTED

	Lakes						Ponds	
	1	2	3	4	5	6	A	B
Total number of taxa	23	131	91	39	72	62	65	43
Sample type:								
Plankton	3	5	4	1	4	3	1	2
Tychoplankton	--	1	1	2	--	3	--	1
Sludge	--	3	3	--	1	1	--	--
Benthic flocculum	1	5	--	1	4	--	1	--
Seepage	--	2	--	--	--	--	--	--
Epilithic	2	1	6	--	--	1	--	--
Total number of samples	6	17	14	4	9	8	2	3
pH	6.0	6.4-6.6	6.6	6.0-6.3	5.8-6.4	5.8	6.0	6.1
Air temperature (°C)	13.3-23.3	10-12.8	20	13.9-16.7	12.2-18.3	18.9	13.3-23.3	--
Water temperature (°C)	14.4-16.7	15.5-16.7	18.9	16.7	15.5-20	15.5	14.4-16.7	--
Aquatic plants mentioned in fieldnotes		Elodea Nitella Isotetes Carex	Elodea Moss Isotetes	Elodea Carex	Elodea Carex	Elodea Ranuncu-lus	Sparganium Callitriche	Elodea
Depth	--	16 m	--	--	--	--	15 cm	--

MATERIALS AND METHODS

COLLECTION

The 63 samples (Table 1) used in this study were collected by Dr. Vinyard during a 9-day helicopter expedition to the Rae Lakes (August 16 - August 25, 1972). Samples were taken from various shoreline positions at each lake, the outlets, some of the connecting creeks, and seepage areas. Open water samples were not collected due to the impracticality of transporting a boat into the area. The number of samples collected from each lake was determined by the variety of habitats encountered. Samples were preserved in 5% formalin solution and returned to the laboratory for examination. A pinch of copper sulfate was added to each liter of preservative to maintain the original colors of the algae.

Each sample was labeled according to the lake and habitat from which it was collected. Habitats sampled included:

1. Planktonic (collected with a #25 plankton net).
2. Tychoplanktonic (squeezing of mosses, Elodea, Nitella, and floating mats of Oedogonium and Rhizoclonium).
3. Sludge.
4. Benthic flocculum.*
5. Seepage.
6. Epilithic.

---

*Vinyard coined this phrase to describe the thin layer of debris often found suspended just off the lake bed in the littoral zone.

PREPARATION AND EXAMINATION

Three permanent slides of each sample were prepared using Karo® syrup with phenol as the mounting medium. Cover slips were sealed with clear fingernail polish. The preserved samples and permanent slides are on file at California State University-Humboldt's cryptogamic herbarium. Each slide was examined and each form, when first encountered, was drawn with the aid of a camera lucida. Wet mounts of most samples were also examined for forms not encountered on the permanent slides and to clarify details overlooked on previously observed plants. Cell dimensions were measured from camera lucida drawings with the aid of a stage micrometer.

An iconographic record was established during preliminary examination of the permanent slides. The illustrations were organized by genus as they were drawn. In this way, accurate records were kept on the occurrence of each plant on a sample-by-sample basis without the immedate need of a species name. Distributions within habitats were easily derived from these records. The systematics presented here follow Prescott (1970) wherever possible. The following occurrence categories were established to indicate species abundance but they are of a subjective nature and should be interpreted in conjunction with the habitat and distribution records; Very rare - seen once or twice, usually in the same sample; Rare - seen several times in one or more samples; Common - low population density, but occurring in many samples; Very common - high population density in one or more samples, or occurring in most samples.

® Registered trademark

RESULTS AND DISCUSSION

Descriptions of species, varieties and forms identified from Rae Lakes and Ponds as well as occurrence and habitat information are given in Appendix A. Species illustrations are presented in Appendix B, while a genus and species index can be found on page 106.

The algal flora of the Rae Lakes were dominated by Chlorophyta (82 percent), most of the remaining taxa were members of Cyanophyta (13 percent). Other divisions were poorly represented except Pyrrhophyta where a single species, Peridinium Willei, occurred quite commonly in large numbers (Table 2). These findings agree with Thomasson's (1962) list of 74 taxa from the Lake Tahoe region, where 85 percent of the taxa were chlorophytes and about 10 percent cyanophytes, and Vinyard's (1951) list of 161 taxa from eight alpine and subalpine regions of the Western United States, where 86 percent were chlorophytes and 13 percent cyanophytes (neither report addressed the diatoms).

Desmids, reflecting the soft water character of the Rae Lakes, were the dominating group within Chlorophyta (65 percent). This relationship also agreed with data collected by Vinyard (1951) and Thomasson (1962) where desmid taxa accounted for 67 and 65 percent of the chlorophytes respectively. Numerous papers have appeared in recent years dealing with algae, especially the desmids, of the North American arctic and subarctic (Whelden 1947, Croasdale 1955, 1956, 1957, 1958, and others). Croasdale (1965) noted that Cosmarium species usually made up more than 50 percent of the arctic desmid flora while some genera (e.g., Micrasterias and Triploceras) were absent. In comparison, 40 percent of the Rae Lakes desmids were Cosmarium, while Micrasterias was represented by eight species. Triploceras was not found in the Rae Lakes.

Non-desmid chlorophytes included 39 chlorococcaleans primarily in the genera Pediastrum, Oocystis, Scenedesmus and Coelastrum. This is apparently the first report of P. taylori since Sieminska described it in 1965 although, as she pointed out, it may have been placed in P. tricornutum var. alpinum by various workers in North America (Taylor 1922, 1924; and Vinyard 1951). Oocystis arctica Prescott is reported here probably for the first time since its description (Prescott and Vinyard 1965).

The seven members of the Euglenophyta reported here were found only in Lake 3. All were in the same sample (sludge near the outlet of Lake 3) except Lepocinclis fusiformis var. major which was in an epilithic sample of Tetraspora lamellosa.

TABLE 2. SUMMARY OF THE TAXA ENCOUNTERED

Divisions	Orders	Families	Genera	Species	Varieties	Forms	Total*
Chlorophyta	9	20	49	119	42	12	173
Euglenophyta	1	1	4	5	2	--	7
Pyrrhophyta	1	1	1	1	--	--	1
Chrysophyta	2	2	2	2	--	--	2
Cyanophyta	3	4	18	24	3	--	27
Total	16	28	74	151	47	12	210

*Total of species, varieties and forms.

Cosmarium askenasyii was considered an Australian and South Asian species until Carter (1935) found and named a variety in British Columbia. The variety was rare in the Rae Lakes material and found only on wet soil. C. subtumidum var. minutum was considered by Krieger and Gerloff (1965) to be a tropical species. It too had a unique microhabitat in a submersed epilithic sample of Tetraspora lamellosa.

Although the study lakes were located in the same small canyon with water from one feeding the next, there were interesting differences in the algal compositions (Table 3). Lake 1, the cirque lake, had the lowest diversity of the study lakes, due primarily to the lack of diverse habitats (e.g., little or no sediment and the absence of vascular aquatic vegetation). Lake 4 also had relatively few taxa compared to the other lakes. It was the smallest of the lakes, very shallow, and nearly overgrown by a Carex marsh. This emergent plant may have acted as a light shield hampering algal growth.

The composition of the major algal groups changed consistently from one end of the chain to the other. There was an increase in blue-green species from 9 percent in Lake 1 to 19 percent in Lake 6 with a corresponding decrease in desmids from 57 percent in Lake 2 to 35 percent in Lake 6. These changes suggested a slight increase in nutrient load down the chain of lakes. Lake 1 did not fit the pattern as desmids comprised 22 percent of the species present.

Although there were numerous desmids present in the Rae Lakes, relatively few were common to all of the lakes and ponds; to the contrary, most were limited to one or two lakes. By comparison, 18 of the 60 non-desmid chlorophytans (30 percent) were found in five or more of the lakes and ponds, whereas only 12 of the 113 desmids (11 percent) were found in five or more of

the lakes and ponds. In addition, 16 taxa (mostly desmids) were limited to pond A with the exception of one Micrasterias also found in pond B. It should be noted that pond A was very shallow and had accumulated sediments, and probably provided a very different habitat from the other lakes and pond B.

The distributions noted in Table 3 may be misleading as the samples were collected during a very short period of time, and it is possible that many of the rarer species become increasingly abundant at different times during the year.

TABLE 3. DISTRIBUTION OF ALGAL SPECIES BY LAKE AND POND

	LAKE						POND		OCCUR.
	1	2	3	4	5	6	A	B	
CHLOROPHYTA									
Chlamydomonadaceae									
Chlamydomonas angulosa					x				VR*
Volvocaceae									
Eudorina elegans	x	x	x		x	x		x	C*
Pandorina morum	x	x			x				R*
Tetrasporaceae									
Tetraspora lamellosa		x	x		x				VC*
Gloeocystaceae									
Asterococcus limneticus		x	x		x	x	x		C
Gloeocystis ampla		x		x	x	x	x		C
G. gigas	x	x	x	x	x	x	x	x	C
G. vesiculosa		x	x		x	x	x	x	C
Chlorococcaceae									
Tetraëdron gracile					x				VR
Palmellaceae									
Sphaerocystis schroeteri	x	x		x	x	x	x	x	C
Oocystaceae									
Ankistrodesmus falcatus		x	x		x		x		C
A. gelifactum		x			x				VR
Eremosphaera viridis				x	x				C
Kirchneriella lunaris var. irregularis		x	x						VR
Oocystis arctica								x	VR
O. borgei							x		R
O. elliptica?					x				R
O. parva	x	x	x	x	x	x	x	x	C
O. pusilla						x			VR

*VR = Very Rare, R = Rare, C = Common, VC = Very Common

(continued)

TABLE 3. DISTRIBUTION OF ALGAL SPECIES BY LAKE AND POND (Continued)

	LAKE						POND		OCCUR.
	1	2	3	4	5	6	A	B	
Oocystaceae (continued)									
O. sp.						x			C
Quadriqula lacustris		x	x	x	x	x	x	x	VC
Zoochlorella spp.		x			x		x		-
Micractiniaceae									
Golenkinia paucispina							x		VR
Dictyosphaeriaceae									
Botryococcus braunii		x	x	x	x	x	x	x	VC
Dictyosphaerium pulchellum	x		x			x			VC
Scenedesmaceae									
Coelastrum cambricum var. intermedium		x							R
C. microporum		x	x		x	x		x	C
C. printzii		x							VR
C. proboscideum			x						VR
Crucigenia rectangularis	x	x	x		x	x	x	x	VC
C. quadrata						x			VR
Enallax alpina		x							VR
Scenedesmus bijuga							x		VR
S. bijuga var. alternans		x							VR
S. dimorphus							x		VR
S. quadricauda	x	x	x		x		x	x	VC
Tetrademus sp.		x							VR
Hydrodictyaceae									
Pediastrum angulosum				x			x		C
P. boryanum		x	x	x	x	x	x	x	VC
P. boryanum var. undulatum		x				x			R
P. braunii			x						VR
P. integrum		x	x						VR
P. muticum		x						x	VR
P. taylori		x					x		R
P. tetras forma		x							R
P. tetras var. tetraodon		x			x	x			C
Sorastrum spinulosum							x		VR

(continued)

TABLE 3. DISTRIBUTION OF ALGAL SPECIES BY LAKE AND POND (Continued)

	LAKE						POND		OCCUR.
	1	2	3	4	5	6	A	B	
Ulotrichaceae									
Ulothrix variabilis		x	x	x	x				C
Chaetophoraceae									
Microthamnion strictissimum	x	x	x	x	x	x		x	C
Protoderma viride								x	C
Coleochaetaceae									
Coleochaete orbicularis				x	x			x	C
Oedogoniaceae									
Bulbochaete sp.	x	x	x	x	x	x	x	x	VC
Oedogonium spp.	x	x	x	x	x	x		x	VC
Cladophoraceae									
Rhizoclonium crassipellitum		x	x	x		x			C
Zygnemataceae									
Mougeotia spp.?	x	x	x		x	x	x	x	C
Spirogyra gracilis	x	x	x		x	x		x	C
Zygnema sp.						x			C
Mesotaeniaceae									
Cylindrocystis brebissonii		x							R
Gonatozygon aculeatum	x		x			x			C
G. brebissonii		x	x		x	x		x	VR
Netrium digitus		x	x		x	x		x	VC
N. digitus var. naegelii			x		x				VR
N. interruptum		x	x		x				VR
Desmidiaceae									
Closterium dianae?		x	x					x	R
C. dianae var. minor?		x	x						R

(continued)

TABLE 3. DISTRIBUTION OF ALGAL SPECIES BY LAKE AND POND (Continued)

	LAKE						POND		OCCUR.
	1	2	3	4	5	6	A	B	
Desmidiaceae (continued)									
C. gracile							x		R
C. intermedium		x	x						C
C. jenneri							x		R
C. libellula			x						VR
C. lunula forma gracilis		x	x	x		x		x	VC
C. macilentum var. japonicum forma					x	x			C
C. rostratum		x			x				R
C. striolatum		x	x		x		x		VC
Cosmarium abbreviatum forma minor			x						VR
C. amoenum		x	x						C
C. angulosum					x			x	R
C. askenasyii var. americana		x							R
C. bioculatum							x		C
C. bipunctatum		x	x	x			x		C
C. botrytis var. tumidum		x	x	x	x	x	x		VC
C. connatum		x			x				VR
C. contractum var. ellipsoideum		x			x				C
C. crenatum			x						C
C. cymatonotophorum var. granulatum		x							VR
C. difficile var. dilatatum		x							R
C. formulosum	x	x	x	x	x	x			C
C. globosum		x	x				x		C
C. hammeri		x							R
C. humile var. striatum				x				x	R
C. impressulum α		x							R
C. impressulum β		x						x	R
C. intermedium			x				x		R
C. laeve var. nov.?				x					R
C. margaritatum forma subrotundata		x							R
C. novae-semliae var. granulatum			x				x		R
C. ornatum var. perornatum			x		x				R

(continued)

TABLE 3. DISTRIBUTION OF ALGAL SPECIES BY LAKE AND POND (Continued)

	LAKE						POND		OCCUR.
	1	2	3	4	5	6	A	B	
Desmidiaceae (continued)									
C. portianum		x	x						C
C. pseudoarctoum					x				R
C. pseudoprotuberans forma minus	x						x		R
C. pseudopyramidatum var. extensum		x							C
C. pseudoquadratulum		x							R
C. pyramidatum		x							C
C. quadratum forma willei			x						C
C. quinarium		x							C
C. rectangulare var. hexagonium		x					x	x	C
C. reniform			x	x		x		x	C
C. sexangulare			x						R
C. sexnotatum var. tristriatum		x	x			x			C
C. subarctoum									
C. suberenatum		x							R
var. isthmochondrum		x	x	x	x	x		x	C
C. subspeciosum var. validius morpha		x					x		C
C. subtumidum var. minutum			x						VR
C. tenue var. depressum		x			x				VR
C. trachypleurum var. fallax				x					VR
C. undulatum var. minutum			x						R
C. ungerianum forma							x		VC
C. venustum var. excavatum		x							R
Desmidium grevillii								x	VR
Euastrum ansatum var. pyxidatum							x		R
E. ansatum var. triporum		x							R
E. bidentatum var. speciosum			x				x		C
E. denticulatum var. angusticeps		x					x		R
E. didelta		x							R
E. elegans					x	x	x		C
E. gemmatum forma latior		x	x						R
E. obesum		x							R
E. oblongum		x					x		C
E. verrucosum var. perforatum		x							R

(continued)

TABLE 3. DISTRIBUTION OF ALGAL SPECIES BY LAKE AND POND (Continued)

	LAKE						POND		OCCUR.
	1	2	3	4	5	6	A	B	
Demidiaceae (continued)									
E. verrucosum var. rhomboideum		x	x	x		x	x		C
E. verrucosum var. vallesiacum		x							R
Micrasterias americana		x							R
M. conferta		x							R
M. denticulata var. angulosa		x		x					R
M. muricata var. tumida							x		VR
M. pinnatifida		x							C
M. rotata							x	x	C
M. rotata forma evoluta		x			x			x	C
M. truncata var. neodamensis		x							R
Penium spirostriolatum			x			x			VR
Pleurotaenium ehrenbergii		x			x	x			VC
P. trabecula var. elongatum		x	x	x	x	x			R
P. trabecula var. maximum		x							VR
Spondylosium planum	x	x	x	x	x	x	x	x	R
S. pulchellum		x							VR
Staurastrum acarides		x							VR
S. alternans		x							C
S. anatinum		x	x		x	x	x	x	C
S. arctiscon	x	x	x	x	x	x		x	VC
S. brebissonii		x	x						R
S. breviaculeatum		x		x			x		VC
S. brevispinum					x			x	R
S. crenulatum						x			VR
S. cuspidatum							x		C
S. dickiei var. circulare		x						x	VR
S. grande var. angulosum		x							C
S. kurilense		x	x						R
S. kurilense forma triquetra							x		VR
S. margaritaceum		x							VR
S. orbiculare			x				x		R
S. pachyrhynchum		x							C
S. polonicum			x						C
S. polymorphum		x	x						R
S. punctulatum		x	x			x	x		C

(continued)

TABLE 3. DISTRIBUTION OF ALGAL SPECIES BY LAKE AND POND (Continued)

	LAKE						POND		OCCUR.
	1	2	3	4	5	6	A	B	
Desmidianceae (continued)									
S. punctulatum var. kjellmani		x							C
S. pyramidatum forma nov.?		x							R
S. sebaldii var. ornatum							x		C
S. spongiosum		x							VR
S. subavicula		x							VR
S. vestitum						x	x		C
Tetmemorus laevis		x	x						VR
Xanthidium subhastiferum	x	x	x	x	x	x	x	x	VC
Characeae									
Nitella sp.		x	x	x	x	x			VC
EUGLENOPHYTA									
Euglenaceae									
Euglena spirogyra var. marchica			x						VR
Lepocinclis fusiformis			x						R
L. fusiformis var. major			x						VR
Phacus sp.			x						VR
Trachelomonus bacillifera var. minima			x						R
T. lacustris			x						R
PYRRHOPHYTA									
Peridiniaceae									
Peridinium willei	x	x	x		x	x	x	x	VC
CHRYSOPHYTA									
Chlorobotrydaceae									
Ducellieria chodatii		x							VR

(continued)

TABLE 3. DISTRIBUTION OF ALGAL SPECIES BY LAKE AND POND (Continued)

	LAKE						POND		OCCUR.
	1	2	3	4	5	6	A	B	
Dinobryaceae									
Dinobryon cylindricum	x				x	x			R
CYANOPHYTA									
Chroococcaceae									
Aphanocapsa elachista var. conferta		x	x	x	x	x	x		C
A. pulchra		x							R
Aphanothece stagnina		x	x		x				C
Chroococcus prescottii			x		x	x			C
C. turgidus		x	x	x	x		x	x	C
Coelosphaerium kuetzingianum						x	x		R
C. pallidum?		x				x	x		C
Dactylococcopsis smithii					x				VR
Eucapsis alpina var. minor							x		R
Gloeothece rupestris?						x			VR
Merismopedia elegans		x			x		x		VC
Microcystis aeruginosa		x		x	x	x	x	x	C
Oscillatoriaceae									
Lyngbya aerugineo-caerulea	x	x			x				R
Oscillatoria agardhii		x	x		x	x			C
O. Limnetica			x						C
O. tenuis							x		C
O. tenuis var. natans		x							C
Phormidium corium			x						R
Nostocaceae									
Anabaena oscillarioides		x	x	x	x	x			VC
A. sphaerica		x			x				C
Cylindrospermum alatosporum					x				R
Nostoc parmelioides			x			x			C
N. paludosum	x	x	x	x	x	x	x	x	VC

(continued)

TABLE 3. DISTRIBUTION OF ALGAL SPECIES BY LAKE AND POND (Continued)

	LAKE						POND		OCCUR.
	1	2	3	4	5	6	A	B	
Scytonemataceae									
Scytonema mirabile		x		x					C
Tolypothrix distorta		x	x	x	x	x	x		C
Rivulariaceae									
Calothrix epiphytica		x							C
C. fusca			x			x			R
CHLOROPHYTA									
desmids	6	75	45	16	25	22	32	19	
non desmid	13	38	26	16	31	26	22	20	
EUGLENOPHYTA	-	-	7	-	-	-	-	-	
PYRRHOPHYTA	1	1	1	-	1	1	1	1	
CHRYSOPHYTA	1	1	-	-	1	1	-	-	
CYANOPHYTA	2	16	12	7	14	12	10	3	
TOTAL	23	131	91	39	72	62	65	43	

LITERATURE CITED

Bourrelly, R. 1966. Les algues d'eau douce. Tome 1. Editions N. Boubee et Cie, Paris. 511 pp.

_____. 1968. Les algues d'eau douce. Tome 2. Editions N. Boubee et Cie, Paris. 438 pp.

Bowman, G. A. 1964. Phytoplankton of the San Joaquin River (exclusive of diatoms). Thesis for M.A. degree, Humboldt State College (unpublished). 147 pp.

Carter, N. 1935. Alpine desmids from British Columbia. Jour. Linn. Soc. Bot. 50(333):151-179.

Coleman, G. A. 1925. Biological survey of the Rae Lakes Region. California Fish & Game. 11(2):57-62.

Croasdale, H. T. 1955. Freshwater algae of Alaska. 1. Some desmids from the interior. Farlowia. 4(4):513-565. Pls. 1-13.

_____. 1956. Freshwater algae of Alaska. 1. Some desmids from the interior. Part 2. Actinotaenium, Micrasterias and Cosmarium. Trans. Am. Microsc. Soc. 75(1):1-70. Pls. 1-17.

_____. 1957. Freshwater algae of Alaska. 1. Some desmids from the interior. Part 3. Cosmarieae concluded. Trans. Am. Microsc. Soc. 76(2):116-158. Pls. 1-10.

_____. 1958. Freshwater algae of Alaska. 2. Some new forms from the plankton of Karluk Lake. Trans. Am. Microsc. Soc. 77(1): 31-35. 1 Pl.

_____. 1965. Desmids of Devon Island, N. W. T., Canada. Trans. Am. Microsc. Soc. 84(3):301-355. Pls. 1-8.

_____. 1973. Freshwater algae of Ellesmere Island, N. W. T. National Museum of Natural Sciences. Publications in Botany. No. 3, Ottawa. 131 pp.

Croasdale, H. and R. Grönblad. 1964. Desmids of Labrador. 1. Desmids of the Southeastern Coastal Area. Trans. Amer. Micros. Soc. 83(2):142-212.

Desikachary, T. V. 1959. Cyanophyta. Academic Press, N. Y. 686 pp.

Grönblad, R. 1920. Finnlandische Desmidiacean aus Keuru. Acta. Soc. Fauna Flora Fenn. 47(4):1-98. Pls. 1-6.

_____. 1962. Desmids from the British Isles. Nova Hedwigia, Band IX: 4(3 and 4):467-479.

Hirano, M. 1968. Desmids of Arctic Alaska. Contrib. Biol. Lab. Kyoto Univ., 21:1-53.

Huber-Pestalozzi, G. 1955. Das Phytoplankton des Süsswassers. Systematik und Biologie. 4. Teil. Stuttgart. 606 pp. Pls. 1-114.

Hylander, C. J. 1928. The algae of Connecticut. Connecticut State Geol. Natur. Hist. Surv. Bull., No. 42. 245 pp.

Irénée-Marie, F. 1952. Contribution a la connaissance des Desmidiees de la region du Lac-St-Jean. Hydrobiologia. 4:1-208. Pls. 1-19.

Jahn, L. J. and F. L. Jahn. 1949. How to know the Protozoa. Wm. C. Brown Co., Dubuque, Iowa. 234 pp. 394 figs.

Johnson, L. P. 1944. Eugleneae of Iowa. Trans. Americ. Microsc. Soc., 63:97-135. Pls. 1-6.

Krieger, W. 1937. Die Desmidiaceen. In L. Rabenhorst, Kryptogamen-Flora von Deutschland, Österreich und der Schweiz. 13, (Abt. 1, Teil 1):377-712. Pls. 1-60.

_____. 1939. Die Desmidaceen. In L. Rabenhorst, Kryptogamen-Flora von Deutschland, Österreich und der Schweiz. 13, (Abt. 1, Teil 2):1-117. Pls. 97-142.

Krieger, W. and J. Gerloff. 1962. Der Gattung Cosmarium. Lf. 1:iii-xvii + 112 pp. Pls. 1-23.

_____. 1965. Der Gattung Cosmarium. Lf. 2:113-240. Pls. 23-42.

Moore, J. G. 1963. Geology of the Mount Pinchot Quadrangle, southern Sierra Nevada, California. Geological Survey Bulletin 1130. 152 pp.

Prescott, G. W. 1962. Algae of the western Great Lakes area. W. C. Brown Co. 977 pp. Pls. 1-136.

_____. 1970. How to know the freshwater algae. W. C. Brown Co. 348 pp.

Prescott, G. W., H. T. Croasdale and W. C. Vinyard. 1972. Desmidiales. 1. Saccodermae, Mesotaeniaceae. N. Am. Flora. II. 6:1-84.

Prescott, G. W., H. T. Croasdale and W. C. Vinyard. 1975. A synopsis of the North American Desmids. Part II. Desmidiaceae: Placodermae. Section 1. University of Nebraska Press, Lincoln, Nebraska. 275 pp.

Prescott, G. W., H. T. Croasdale and W. C. Vinyard. 1977. A synopsis of the North American Desmids. Part II. Desmidiaceae: Placodermae. Section 2. University of Nebraska Press, Lincoln, Nebraska. 413 pp.

Prescott, G. W. and A. M. Scott. 1952. Some south Australian Desmids. Trans. Roy. Soc. S. Aust. 75:55-69.

Prescott, G. W. and W. C. Vinyard. 1965. Ecology of Alaskan freshwater algae V. Limnology and flora of Malikpuk Lake. Trans. Amer. Microsc. Soc. 84(1):98-126.

Scott, A. M. and G. W. Prescott. 1958. Some freshwater algae from Arnhem Land in the northern territory of Australia. Records of the American-Australian Scientific Expedition to Arnhem Land. 3:8-135.

Sieminska, J. 1965. Algae from Mission Wells Pond, Montana. Trans. Amer. Microsc. Soc. 84(1):98-126.

Skuja, H. 1949. Zur Süsswasserflora Burmas. Nova Acta Reg. Soc. Sci. Upsaliensis. Ser. 4, 14(5):1-188, 37 Pls.

Smith, G. M. 1920. Phytoplankton of the inland lakes of Wisconsin. 1. Bull. Wisconsin Geol. and Nat. Hist. Surv. 57:1-243. Pls. 1-51.

_____. 1924. Phytoplankton of the inland lakes of Wisconsin. 2. Bull. Wisconsin Geol. and Nat. Hist. Surv. 57:1-227. Pls. 1-38.

Taylor, W. R. 1922. Notes on some algae from British Columbia. Rhodora. 24:101-111.

_____. 1924. Further notes on British Columbia algae. Rhodora. 26:160-166.

_____. 1935. The freshwater algae of Newfoundland. II. Papers Mich Acad. Sci., Arts, Lett. 20:185-230. Pls. 1-17.

Teiling, E. 1954. Actinotaenium genus desmdiacearum resuscitatum. Botaniska Notiser. Häfte 4:376-426.

Teiling, E. 1967. The desmid genus Staurodesmus. Ark. F. Bot. 6(11):466-629. Pls. 1-30.

Thomasson, C. 1962. Planktological notes from western North America. Ark. F. Bot. 4(14):437-463.

Vinyard, W. C. 1951. Distribution of alpine and subalpine algae in western United States. Thesis for M.S. degree, Michigan State College (unpublished). 55 p. Pls. 1-16.

Wade, W. E. 1957. Additions to our knowledge of the desmid flora of Michigan. Rev. Algol. 4:249-273. Pls. 1,2.

Wailes, C. H. 1930. Protozoa and algae, Mount Ferguson, B. C. Vancouver Mus. and Art Notes. 5:160-165. 1 Pl.

West, W. and G. S. West, 1904. A monograph of the British Desmidiaceae. I. The Ray Society, London. 224 pp. Pls. 1-32.

_____. 1905. A monograph of the British Desmidiaceae. II. 204 pp. Pls. 33-64.

_____. 1908. A monograph of the British Desmidiaceae. III. 273 pp. Pls. 96-128.

_____. 1912. A monograph of the British Desmidiaceae. IV. 191 pp. Pls. 96-128.

West, W., G. S. West and N. Carter. 1923. A monograph of the British Desmidiaceae. V. 300 pp. Pls. 129-167.

Wheldon, R. M. 1947. Botany of the Canadian eastern Arctic. Part II, Thallophyta and Bryophyta. 1. Algae. Nat. Mus. Canada Bull. 97:13-137. Pls. 1-8.

Wolle, F. 1887. Desmids of the Pacific coast. Bull. California Acad. Sci. 2(7):432-437.

APPENDIX A

SPECIES, VARIETY, AND FORM DESCRIPTIONS

The species descriptions are organized as follows:

1. Species, author and date. [An asterisk (*) before a species name denotes a species thought to be a new record for California while a question mark (?) immediately following a portion of the name denotes questionable identity of that part.]

2. Primary reference(s) used in making the identification.

3. Dimensions and descriptive information when it varies from the primary reference. All dimensions are given in micrometers (μm). Parentheses surrounding cell dimensions indicate the occurrence of extremely large or small cells in the sample material. Dimensions for desmids are given with the following abbreviations (applies to single cells):

   L. - length
   W. - maximum width
   Wa. - maximum width at apex
   Th. - maximum thickness
   Tha. - maximum thickness at apex
   Isth. - width of isthmus in face view

4. Occurrence. These data are of a subjective nature and should be interpreted in conjunction with the habitat and distribution records. Categories:

   Very rare - seen once or twice, usually in the same sample.

   Rare - seen several times in one or more samples.

   Common - low population density, but occurring in many samples.

   Very common - high population density in one or more samples, or occurring in most samples.

5. Habitat(s).

6. Appendix B plate and figure numbers with corresponding species illustrations.

SYSTEMATICS

CHLOROPHYTA

Volvocales

Chlamydomonadaceae

Chlamydomonas Ehrenberg

Chlamydomonas angulosa Dill. 1895.
Prescott. 1962: 70, Pl. 1, Fig. 3.
Cells 20 µm long, 13 µm in diameter. This determination was based on two cells which had lost their flagella. In all other respects they closely followed Prescott's description.
Very rare.
Sludge.
Pl. 1, Fig. 2.

Volvocaceae

Eudorina Ehrenberg

Eudorina elegans Ehrenberg 1832.
Prescott. 1962: 76, Pl. 1, Figs. 24-26.
Cells 8-19 µm in diameter. Colonies were up to 165 µm in diameter.
Common in net tows.
Plankton, tychoplankton, benthic flocculum, and epilithic.
Pl. 1, Fig. 1.

Pandorina Bory

Pandorina morum (Müll.) Bory 1824.
Prescott. 1962: 75, Pl. 1, Fig. 23.
Cells 7.5-12 µm long, 5-8 µm in diameter. The colonies were 16-celled; the cells were smaller than Prescott's.
Rare.
Plankton, benthic flocculum, and sludge.
Pl. 1, Fig. 3.

Tetrasporales

Tetrasporaceae

Tetraspora Link

Tetraspora lamellosa Prescott 1944.
Prescott. 1962: 88, Pl. 5, Fig. 6.

Cells 5-7.5 µm in diameter. The cells were smaller than Prescott's but the distinguishing characteristics were present: pseudocilia were greater than 20 times the cell diameter and possessed distinct lamellate cell sheaths. Shape of colony was unknown.
Very common.
Epilithic.
Pl. 1, Fig. 4.

Gloeocystaceae

Asterococcus Scherffel

*Asterococcus limneticus G.M. Smith 1918.
Prescott. 1962: 86, Pl. 4, Fig. 11.
Cells 6-9 µm in diameter, colonies 30-50 µm in diameter. Dimensions of cells and colonies were at the lower end of the range given by Prescott.
Common.
Plankton, tychoplankton, and benthic flocclulum.
Pl. 1, Fig. 5.

Gloeocystis Nägeli

Gloeocystis ampla (Kütz.) Lagerheim 1883.
Prescott. 1962: 84, Pl. 3, Fig. 17.
Cells 10-15 µm long, 7-10 µm in diameter.
Common.
Plankton, tychoplankton, benthic flocculum, and seepage.
Pl. 1, Fig. 11.

Gloeocystis gigas (Kütz.) Lagerheim 1883.
Smith. 1920: 101, Pl. 19, Fig. 2.
Cells 12-18-(20) µm in diameter. Lamellations were rarely obvious.
Common.
Plankton tychoplankton, wet soil, benthic flocculum, and sludge.
Pl. 1, Fig. 10.

Gloeocystis vesiculosa Nägeli 1849.
Prescott. 1962: 85, Pl. 3, Fig. 15.
Cells 3.5-11 µm in diameter.
Common.
Plankton, tychoplankton, benthic flocculum, sludge, and epilithic.
Pl. 1, Fig. 7.

Chlorococcales

Chlorococcaceae

Tetraedron Kützing

*Tetraëdron gracile (Reinsch) Hansgirg 1889 forma.
Prescott. 1962: 265, Pl. 60, Fig. 1.
Cells 11-15 µm in diameter without processes, 30-35 µm in diameter with

processes. They did not have short spines tipping the processes. The illustration was identical to Skuja's (1949; Pl. 10, Figs. 19 and 20), which was named Isthmochloron gracile (Reinsch) Skuja.
Plankton.
Pl. 1, Fig. 6.

Palmellaceae

Sphaerocystis Chodat

Sphaerocystis Schroeteri Chodat 1897.
Prescott. 1962: 83, Pl. 3, Figs. 6 and 7.
Cells (4)-6-22 µm in diameter.
Common.
Plankton.
Pl. 1, Fig. 12.

Oocystaceae

Ankistrodesmus Corda

Ankistrodesmus falcatus (Corda) Ralfs 1848.
Prescott. 1962: 253, Pl. 56, Figs. 5 and 6.
Cells 40-68 µm long, 2.5 µm in diameter.
Common.
Net tow in 15 centimeters of water, benthic flocculum, seepage, and tychoplankton.
Pl. 1, Fig. 9.

*Ankistrodesmus gelifactum (Chod.) Bourrelly 1951.
Croasdale. 1973: 60, Pl. 9, Figs. 32 and 33.
Cells 35 µm long, 6 µm in diameter, single or in pairs.
Very rare.
Wet soil (seepage into Lake 2), and plankton.
Pl. 1, Fig. 8.

Eremosphaera de Bary

*Eremosphaera viridis de Bary 1858.
Prescott. 1962: 240, Pl. 53, Fig. 22.
Cells about 135 µm in diameter.
Common.
Sludge and benthic flocculum.
Pl. 2, Fig. 1.

Kirchneriella Schmidle

Kirchneriella lunaris var. irregularis G. M. Smith 1920.
Cells 8-11 µm long, 4 µm in diameter.
Very rare.
Sludge.

Pl. 2, Fig. 10.

Oocystis Nägeli

The accurate cytoplasmic data needed to make determinations in this genus were not always available leaving several species unidentified. Living material would have been particularly valuable for this genus.

*Oocystis arctica Prescott 1965.
Prescott and Vinyard. 1965: 455, Pl. 8, Figs. 13 and 14.
Cells 11.5-18 µm long, 8-10 µm in diameter. They were smaller than Prescott and Vinyard's which were 19-20 µm by 14-15 µm.
Very rare.
Squeezings of Elodea.
Pl. 2, Fig. 2.

*Oocystis borgei Snow 1903.
Prescott. 1962: 243, Pl. 51, Fig. 10.
Cells 11.5-14 µm long, 10 µm in diameter.
Rare.
Benthic flocculum.
Pl. 2, Fig. 4.

*Oocystis elliptica? W. West 1892.
Prescott. 1962: 244, Pl. 51, Fig. 11.
Cells 27-28 µm long, 13-14 µm in diameter. They were slightly longer than Prescott's and each chloroplast had a pyrenoid. Prescott wrote, "apparently without pyrenoids."
Rare.
Plankton.
Pl. 2, Figs. 5 and 6.

*Oocystis parva West and West 1898.
Prescott. 1962: 246, Pl. 54, Fig. 3.
Cells 9-15-(17) µm long, 4-8-(9) µm in diameter.
Common.
Plankton, tychoplankton, epilithic, and benthic flocculum.
Pl. 2, Fig. 3.

*Oocystis pusilla? Hansgirg 1890.
Prescott. 1962: 246, Pl. 51, Fig. 15; Pl. 54, Figs. 4 and 5.
Cells 8-9 µm long, 4-5 µm in diameter. The single chloroplast was a parietal plate with one pyrenoid. They differed from Prescott's description in the extension of the mother cell wall to form two bluntly rounded poles. Polar nodules were lacking.
Very rare.
Plankton.
Pl. 2, Fig. 7.

Oocystis sp.
Cells 37-40 µm long, 19-25 µm in diameter. Chloroplasts, 14-20 parietal disks, usually with one pyrenoid apiece. Cells were solitary or in groups of

4-8 enclosed in the old mother cell wall. Poles of cells had nodular thickenings.
Common.
Tychoplankton.
Pl. 2, Fig. 8.

Quadrigula Printz

Quadrigula lacustris (Chod.) G. M. Smith 1920.
Prescott. 1962: 260, Pl. 59, Figs. 4 and 5.
Cells 21-23 µm long, 2-3 µm in diameter.
Very common.
Plankton and tychoplankton.
Pl. 2, Fig. 12.

Zoochlorella spp. Zoochlorella Brand
Prescott. 1962: 235.
Zoochlorella was found inhabiting 2 protozoans: Paramecium bursaria, (Jahn. 1949: 187, Fig. 298-D) and Acanthocystis turfacea Leidy, (Jahn. 1949: 104, Fig. 164).

Micractiniaceae

Golenkinia Chodat

*Golenkinia paucispina West and West 1902.
Prescott. 1962: 213, Pl. 45, Fig. 2.
Cells 20 µm in diameter (- spines), spines 10-12 µm long.
Very rare.
Benthic flocculum and sludge.
Pl. 2, Fig. 9.

Dictyosphaeriaceae

Botryococcus Kützing

Botryococcus braunii Kützing 1849.
Prescott. 1962: 232, Pl. 52, Figs. 1,2, and 11.
Cells 6-8 µm long, 3.2-4 µm in diameter. Colonies were up to 200 µm in diameter.
Very common.
Plankton, benthic floccculum, tychoplankton, epilithic, and sludge.
Pl 2, Fig. 13.

Dictyosphaerium Nägeli

Dictyosphaerium pulchellum Wood 1874.
Prescott. 1962: 238, Pl. 51, Figs. 5-7.
Cells 4-9 µm in diameter.

Very common.
Plankton.
Pl. 2, Fig. 11.

Scenedesmaceae

Coelastrum Nägeli

Coelastrum cambricum var. intermedium (Bohlin) G. S. West.
G. W. Prescott (personal correspondence).
Cells 7-9 µm in diameter. Coenobia was about 25 µm in diameter.
Rare.
Benthic flocculum on gravel.
Pl. 3, Fig. 1.

*Coelastrum microporum Näegeli 1855.
Prescott. 1962: 230, Pl. 53, Fig. 3.
Cells 6-13 µm in diameter. Coenobia were occasionally adhering to one another forming compound colonies. The intercellular spaces characteristic of the species were often lacking.
Common.
Plankton, tychoplankton, benthic flocculum, and sludge.
Pl. 3, Figs. 2 and 3.

*Coelastrum printzii Rayss 1915.
Croasdale. 1973: 56, Pl. 8, Figs. 10-12.
Cells 12-14 µm in diameter. They had small triangular intercellular spaces (not shown on Croasdale's illustrations).
Very rare.
Sludge.
Pl. 3, Fig. 4.

*Coelastrum proboscideum Bohlin 1897.
Prescott. 1962: 230, Pl. 53, Figs. 4, 5, and 8.
Cells 10 µm in length, 12 µm in diameter. Coenobium 25 µm in diameter.
Very rare.
Sludge.
Pl. 3, Fig. 7.

Crucigenia Morren

Crucigenia rectangularis (Braun) Gay 1891.
Prescott. 1962: 285, Pl. 65, Figs. 7 and 8.
Cells 6-8 µm long, 3.5-4 µm in diameter.
Very common.
Plankton, tychoplankton, benthic flocculum, sludge, and epilithic.
Pl. 3, Fig. 6.

Crucigenia quadrata Morren 1830 forma.

Prescott. 1962: 285, Pl. 65, Fig. 10.
Cells 8-10 µm long, 8-9 µm in diameter. Differed from type in larger dimensions, about twice the size of other records.
Very rare.
Tychoplankton.
Pl. 3, Fig. 5.

Enallax Pascher

*Enallax alpina Pascher 1943.
Bourrelly. 1966: 220, Pl. 36, Fig. 19
Cells 19-20 µm long, 10-11 µm in diameter, with 4 longitudinal ribs. Coenobium was a cluster of 4 staggered cells.
Very rare.
Sludge.
Pl. 3, Fig. 8.

Scenedesmus Meyen

Scenedesmus bijuga (Turp.) Lagerheim 1893.
Prescott. 1962: 276, Pl. 63, Figs. 2 and 7.
Cells 11-14 µm long, 3-5 µm in diameter.
Very rare.
Benthic flocculum.
Pl. 3, Fig. 22.

Scenedesmus bijuga var. alternans (Reinsch) Hansgirg 1888.
Prescott. 1962: 277, Pl. 63, Figs. 3 and 4.
Cells 10 µm long, 7 µm in diameter.
Very rare.
Sludge.
Pl. 3, Fig. 9.

Scenedesmus dimorphus (Turp.) Kützing 1833.
Prescott. 1962: 277, Pl. 63, Figs. 8 and 9.
Cells 10-16 µm long, 3 µm in diameter.
Very rare.
Benthic flocculum.
Pl. 3, Fig. 11.

Scenedesmus quadricauda (Turp.) Brébisson 1835.
Prescott. 1962: 280, Pl. 64, Fig. 2.
Cells 14-19 µm long, 4-7 µm in diameter. Colonies were 4 and 8-celled.
Very common.
Plankton, tychoplankton, benthic flocculum, and sludge.
Pl. 3, Fig. 10.

Tetradesmus Smith

Tetradesmus sp.
Prescott. 1962: 283, Pl. 64, Figs. 12-14.
Bowman. 1964: 43, Pl. 7, Fig. 1.
Cells 11 µm long, 3-4 µm in diameter. They differed from T. Wisconsinense by the cell ends being extended into sharp spines. This feature was mentioned by Bowman (1964).
Very rare.
Benthic flocculum.
Pl. 3, Fig. 21.

Hydrodictyaceae

Pediastrum Meyen

*Pediastrum angulosum (Ehrenb.) Meneghini.
Prescott and Vinyard. 1965: 449, Pl. 5, Figs. 9-18.
Cells (8)-12-23 µm long, (7)-10-18 µm wide. Eight and 16-celled coenobia had smooth cell walls, while 32-celled coenobia had granulated cell walls.
Common.
Benthic flocculum and tychoplankton.
Pl. 3, Fig. 15, and Pl. 4, Fig. 2.

Pediastrum boryanum (Turp.) Meneghini 1840.
Prescott. 1962: 222, Pl. 47, Fig. 9, and Pl. 48, Figs. 1 and 3.
Cells (10)-13-20-(25) µm long, (7)-10-16 µm wide. Coenobia were 8, 16, 32, and 64-celled. Processes were variable in length and shape.
Very common.
Tychoplankton, sludge, and benthic flocculum.
Pl. 3, Figs. 18, 19, and 23.

*Pediastrum boryanum var. undulatum Wille 1879.
Prescott. 1962: 223, Pl. 48, Fig 2.
Cells 12-28 µm long, 15-20 µm wide.
Rare.
Tychoplankton and benthic flocculum.
Pl. 4, Figs. 1 and 3.

*Pediastrum braunii Wartmann 1862.
Prescott. 1962: 223, Pl. 48, Fig. 5.
Calls 8-12 µm in diameter, smooth walls.
Very rare.
Sludge.
Pl. 3, Fig. 12.

*Pediastrum integrum Nägeli 1849.
Carter. 1973: 55, Pl. 8, Fig. 4.
Prescott. 1962: 225, Pl. 48, Figs. 9 and 10.
Cells 15-25 µm in diameter.
Very rare.

Sludge.
Pl. 3, Fig. 16.

Pediastrum muticum Kützing 1849.
Prescott. 1962: 225, Pl. 49, Fig. 8.
Cells 13-14 µm in diameter. The cells were smaller than Prescott's which were 20 µm in diameter.
Very rare.
Plankton.
Pl. 3, Fig. 13.

*Pediastrum taylori Sieminska 1965.
Sieminska. 1965: 100, Pl. 2, Figs. 8-14.
Cells 11-15 µm long, 12-14 µm in diameter. The entire cell wall was punctate. Colonies were 4, 8, and 16-celled.
Rare.
Tychoplankton, benthic flocculum, and sludge.
Pl. 3, Fig. 17.

Pediastrum tetras (Ehrenb.) Ralfs 1844 forma.
Prescott. 1962: 227, Pl. 50, Figs. 3 and 6.
Smith. 1920: 173, Pl. 48, Figs. 9-12.
Cells 8-10 µm long, 9-10 µm in diameter. They differed from the type in shallower peripheral incision and the margins were straight or only slightly concave. Colonies 8-celled.
Rare.
Tychoplankton and benthic flocculum.
Pl. 3, Fig. 14.

Pediastrum tetras var. tetraodon (Corda) Rabenhorst 1868.
Prescott. 1962: 227, Pl. 50, Fig. 7.
Cells (4.5-6)-8.5-10 µm long, (4.8-5.5)-10-13 µm in diameter. The cells were much smaller than Prescott's.
Common.
Tychoplankton.
Pl. 3, Figs. 20 and 24.

Sorastrum Kützing

Sorastrum spinulosum Nägeli 1849.
Prescott. 1962: 228, Pl. 50, Fig. 9, and Pl. 53, Fig. 1.
Cells 6-7 µm long, 8 µm wide, and 5 µm thick, with short spines, 3 µm long.
Very rare.
Benthic flocculum.
Pl. 4, Fig. 12.

Ulotrichales

Ulotrichaceae

Ulothrix Kützing

*Ulothrix variabilis Kützing 1849.
Prescott. 1962: 97, Pl. 6, Fig. 13.
Cells 11-14 µm long, 6-7 µm in diameter. They had 1, sometimes 2 pyrenoids in each chloroplast.
Common.
Plankton and tychoplankton (was not found attached to a substrate).
Pl. 4, Fig. 18.

Chaetophorales

Chaetophoraceae

Microthamnion Nägeli

*Microthamnion strictissimum Rabenhorst 1859.
Prescott. 1962: 122, Pl. 11, Figs. 5 and 6.
Cells 7-18 µm long, 3-4 µm in diameter.
Common.
Plankton, tychoplankton, benthic flocculum, and sludge (this plant was not found attached in our samples).
Pl. 4, Fig. 7.

Protoderma Kützing

*Protoderma viride Kützing 1843.
Prescott. 1962: 123, Pl. 9, Fig. 10, and Pl. 14, Fig. 10.
Cells 8-10 µm long, 4-5 µm in diameter.
Common.
Epiphytic on Elodea.
Pl. 4, Figs. 8 and 9.

Coleochaetaceae

Coleochaete Brébisson

*Coleochaete orbicularis Pringsheim 1860.
Prescott. 1962: 129, Pl. 18, Figs. 3-5.
Cells about 20 µm long, 10 µm in diameter, and thallus up to 120 µm in diameter.
Common.
Epiphytic on Elodea and Rhizoclonium crassipellitum.
Pl. 4, Fig. 5.

Oedogoniales

Oedogoniaceae

Bulbochaete Agardh

Bulbochaete sp.
Plants were observed only in the vegetative condition.
Very common.
Fragments were found in every habitat.

Oedogonium Link

Oedogonium spp.?
The lack of adequate reproductive structures left species identification(s) in doubt.
Very common in most habitats.

Siphonocladales

Cladophoraceae

Rhizoclonium Kützing

*Rhizoclonium crassipellitum West and West 1897.
Prescott. 1962: 141, Pl. 23, Fig. 1.
Cells (80)-125-185 µm long, 40-50 µm in diameter. Cell walls 5-9 µm thick. The alga did not attain Prescott's larger dimensions and it was in soft water, whereas Prescott reported it in hard water lakes.
Common.
Tychoplankton and as a floating mat (6 x 8 meters).
Pl. 4, Fig. 13.

Zygnematales

Zygnemataceae

Mougeotia Agardh

Mougeotia spp.
Lack of mature reproductive structures made these plants impossible to identify to species.
Common.

Spirogyra Link

Spirogyra gracilis (Hass.) Kützing 1849.
Prescott. 1962: 315, (not illustrated).
Hylander. 1928: 109, Pl. 17, Figs. 12 and 13.
Cells 50-150 µm long, 28-30 µm in diameter. Zygospore 42-44 µm long, 23-25 µm in diameter. They closely followed Prescott's description except the cells had a greater diameter, and the zygospores had a slightly smaller

diameter. Reproductive structures were rare.
Common.
Plankton.
Pl. 5, Figs. 13 and 14.

Zygnema Agardh

Zygnema spp.
They lacked the reproductive structures necessary for species determination.
Common.
Within a floating mat of Oedogonium sp.

Mesotaeniaceae

Cylindrocystis Meneghini

Cylindrocystis brebissonii Meneghini 1838.
Prescott, Croasdale, and Vinyard. 1972: 20, Pl. 2, Figs. 1-5.
L. 32-43 µm, W. 14-15 µm.
Rare.
Sludge.
Pl. 18, Figs. 4 and 5.

Gonatozygon de Bary

*Gonatozygon aculeatum Hastings 1892.
Prescott, Croasdale, and Vinyard. 1972: 33, Pl. 8, Figs. 13-15.
L. 150-269 µm, W. 10-20 µm, Wa. 11-17 µm, Spines 6-6.5 µm.
Common.
Plankton, squeezings of Ranunculus, and sludge.
Pl. 5, Figs. 6 and 7.

*Gonatozygon brebissonii de Bary 1858.
Prescott, Croasdale, and Vinyard. 1972: 34, Pl. 8, Figs. 1-3 and 11.
L. 170-190 µm, W. 7.5-9 µm, Wa. 6 µm.
Very rare.
Benthic flocculum on gravel and tychoplankton (with Oedogonium).
Pl. 5, Fig. 5.

Netrium Nägeli

Netrium digitus (Ehrenb.) Itzigsohn and Roth 1856.
Prescott, Croasdale, and Vinyard. 1972: 24, Pl. 4, Figs. 14-16 and 21.
L. 142-312 µm, W. 42-82 µm.
Very common.
Sludge, seepage, and plankton.
Pl. 6, Fig. 15.

Netrium digitus var. naegelii (Bréb.) Krieger 1935.
Prescott, Croasdale, and Vinyard. 1972: 26, Pl. 5, Figs. 3 and 4.
L. 123 µm, W. 28 µm.
Very rare.
Sludge.
Pl. 6, Fig. 14.

*Netrium interruptum (Bréb.) Lütkemüller 1902.
Prescott, Croasdale, and Vinyard. 1972: 28, Pl. 5, Figs. 17 and 18.
L. 178 µm, W. 40 µm.
Very rare.
Sludge.
Pl. 6, Fig. 16.

Desmidiceae

Closterium Nitzsch

Closterium dianae? Ehrenberg 1838.
Prescott, Croasdale, and Vinyard. 1975: 46, Pl. 23, Figs. 16, 16a, and 17.
L. 192-2-214 µm, W. 20-21 µm, Wa. 3.5-5 µm.
Rare.
Tychoplankton.
Pl. 6, Fig. 11.

Closterium dianae var. minor? Hieronymus 1895.
Prescott, Croasdale, and Vinyard. 1975: 47, Pl. 23, Figs. 8 and 12.
L. 110-154 µm, W. 17.5-25 µm, Wa. 3-6 µm.
Rare.
Seepage and with Tetraspora lamellosa.
Pl. 6, Fig. 12.

Closterium gracile Brébisson 1839.
Prescott, Croasdale, and Vinyard. 1975: 52, Pl. 16, Figs. 2, 15, and 16.
L. 186-208 µm, W. 7-8 µm, Wa. 2.5-3 µm.
Benthic flocculum.
Rare.
Pl. 6, Figs. 3 and 4.

Closterium intermedium Ralfs 1848.
Prescott, Croasdale, and Vinyard. 1975: 54, Pl. 29, Figs. 10, 10a, and 11.
L. 245-275 µm, W. 15-16 µm, Wa. 6-7 µm.
Common.
Benthic flocculum and sludge.
Pl. 6, Fig. 10.

Closterium jenneri Ralfs 1848.
Prescott, Croasdale, and Vinyard. 1975: 55, Pl. 23, Figs. 4 and 10.
L. 69-100 µm, W. 10-15 µm, Wa. 2.5-3 µm.
Rare.
Benthic flocculum and sludge.
Pl. 6, Fig. 13.

Closterium libellula Focke 1847.
Prescott, Croasdale, and Vinyard. 1975: 60, Pl. 12, Fig. 12.
L. 242 µm, W. 38 µm, Wa. 16 µm.
Very rare.
Sludge.
Pl. 6, Fig. 1.

Closterium lunula forma gracilis Messikommer 1935.
Prescott, Croasdale, and Vinyard. 1975: 66, Pl. 14, Fig. 15.
L. 420-590 µm, W. 80-100 µm, Wa. 20-21.5 µm.
Very common.
Benthic flocculum, tychoplankton, and sludge.
Pl. 6, Fig. 9.

Closterium macilentum var. japonicum (Sur.) Grönblad 1926 forma.
Prescott, Croasdale, and Vinyard. 1975: 68, Pl. 22, Fig. 11.
L. 480-550 µm, W. 35-40 µm, Wa. 7-9 µm.
Common.
Plankton and sludge.
Pl. 6, Figs. 5 and 6.

Closterium rostratum Ehrenberg 1832.
Prescott, Croasdale, and Vinyard. 1975: 83, Pl, 31, Figs. 3 and 12.
L. 312 µm, W. 21 µm, Wa. 4 µm.
Rare.
Benthic flocculum on gravel shore.
Pl. 6, Figs. 7 and 8.

Closterium striolatum Ehrenberg 1832.
Prescott, Croasdale, and Vinyard. 1975: 87, Pl. 27, Figs. 1, 3, 10 (fa.), and 14, and Pl. 28, Fig. 4.
L. 247-320 µm, W. 30-55 µm, Wa. 10-22 µm. An unusual form of the species was also found in sludge sample 17.14 from Lake 2 containing numerous Closterium identical to C. striolatum except for the large numbers of axial pyrenoids, 9-25 per chloroplast.
Very common.
Sludge and benthic flocculum.
Pl. 6, Fig. 2.

Cosmarium Corda

Many species of Cosmarium were transferred to the genus Actinotaenium by Teiling (1954). Since his nomenclature has been accepted by many students of the desmids, both names, where applicable, are given in this text.

*Cosmarium abbreviatum Racib. forma minor West and West 1908.
West and West. 1908: 85, Pl. 72, Fig. 12.
L. 10 µm, W. 12 µm, Th. 5.5 µm, Isth. 4 µm. It was slightly larger than the Wests'. Krieger and Gerloff (1965), showed a name change to C. abbreviatum Racib. forma minus along with their illustration, but did not include a description (this is to follow in a forthcoming Part 3).
Very rare.
Sludge.
Pl. 7, Fig. 10.

Cosmarium amoenum Brébisson 1849.
West and West. 1912: 29, Pl. 102, Figs. 1-4, and Pl. 103, Fig. 9.
L. 57-58 µm, W. 31-33 µm, Th. 30 µm, Isth. 16-18 µm.
Common.
Moss squeezings and wet soil.
Pl. 10, Fig. 2.

*Cosmarium angulosum Brébisson 1856.
Krieger and Gerloff. 1965: 190, Pl. 37, Fig. 13.
L. 17 µm, W. 12.5 µm, Isth. 4 µm.
Rare.
Squeezings of Elodea and benthic flocculum.
Pl. 8, Fig. 14.

*Cosmarium askenasyii Schmidle var. americana Carter 1935.
Carter. 1935: 160, Figs. 43-45.
L. 82-92 µm, W. 66-74 µm, Th. 43 µm, Isth. 24-28 µm.
Rare.
Wet soil.
Pl. 9, Fig. 10.

Cosmarium bioculatum Brébisson 1835.
West and West. 1905: 165, Pl. 61, Figs. 3-7.
L. 24-25 µm, W. 18-20 µm, Th. 11-13 µm, Isth. 6.5-7 µm. They were slightly larger than the Wests'.
Common.
Net tow through 15 centimeters of water. Note: Most of the numerous desmids in this net tow sample had strands of mucilage extending out from their pores as illustrated on Pl. 8, Fig. 12.
Pl. 8, Figs. 7 and 12.

*Cosmarium bipunctatum Börge. 1890.
West and West. 1908: 213, Pl. 85, Fig. 6.
L. 22-24 µm, W. 19-22 µm, Th. 15-16 µm, Isth. 6-8 µm. Ours was longer than wide.
Common.
Benthic flocculum and epilithic with Tetraspora lamellosa.
Pl. 11, Fig. 8.

Cosmarium botrytis var. tumidum Wolle 1892. forma?
There are noteworthy variations in several reports of this plant

including my own. Wolle's plant, as originally described, apparently had a central swelling on the face of the semicell and larger granules in the central area. West and West (1912) noted, "often with one large granule adjacent to the isthmus", otherwise similar to Wolle's plant. Croasdale's (1956) plant lacked the central swelling but had a basal granule. Wade (1957) formed a new taxon, C. botrytis var. tumidum forma nudum, which was similar to Croasdale's plant and mine. Not having seen Wolle or Wade's descriptions I hesitate to assign this plant to a specific taxon. The following description will suffice until more information becomes available.

L. 70-77 µm, W. 55-62 µm, Th. 34-36 µm, Isth. 13-16-(19) µm. The semicells were semi-circular, with slightly flattened apices, were circular to broadly oval in side view, and were elliptic with sides parallel in end view. Granules were in more or less concentric series within the margin, becoming larger and scattered in the central area. The cell wall lacked granules on the apex and on both sides of the central area, and was punctate everywhere except smooth areas on the face of the semicell. The basal granule was always present.

Very common.
Found in most habitats.
Pl. 9, Fig. 3.

*Cosmarium connatum Brebisson 1848.
West and West. 1908: 25, Pl. 67, Figs. 15-17.
L. 73 µm, W. 51 µm, Isth. 38 µm.
Very rare.
Benthic flocculum.
Pl. 11, Fig. 11.

*Cosmarium contractum var. ellipsoideum (Elfv.) West and West 1905.
West and West. 1905: 172, Pl. 61, Figs. 28 and 35.
Prescott and Scott. 1952: 8, Fig. 3, No. 8.
L. 36-40 µm, W. 29-32 µm, Th. 21 µm, Isth. 9.5-11 µm. Wall was delicately punctate.
Common.
Net tow through 15 centimeters of water and sludge.
Pl. 8, Fig. 8.

Cosmarium crenatum Ralfs 1844.
West and West. 1912: 35, Pl. 98, Figs. 9-12.
L. 36 µm, W. 32 µm, Th. 23 µm, Isth. 7 µm.
Common.
Sludge.
Pl. 10, Fig. 5.

*Cosmarium cymatonotophorum W. West var. granulatum Grönblad.
G. W. Prescott (personal correspondence).
L. 15 µm, W. 14 µm, Th. 12 µm, Isth. 7.5 µm. The central protuberance was more pronounced than in the type. The granules were flat and difficult to see.
Very rare.
Sludge.

Pl. 11, Fig. 6.

*Cosmarium difficile Lütkem. var. dilatatum Börge forma nov.
G. W. Prescott (personal correspondence).
L. 25-29 µm, W. 16-18 µm, Th. 10-11 µm, Isth. 4.5 µm. They lacked the two transverse series of wall pits characteristic of the variety. Wall had minute, dense punctae.
Rare.
Sludge.
Pl. 8, Figs. 1 and 2.

Cosmarium formulosum Hoff 1888.
West and West. 1908: 240, Pl. 88, Figs. 1-3.
L. 40-43 µm, W. 35-39 µm, Th. 22 µm, Isth. 10-14 µm. Granulation was variable due to frequent fusion of granules. Central protuberance had 3-5 vertical rows of granules, subtended by a curved row of 5 granules. Punctae were sometimes present between the granules on the central protuberance.
Common.
Squeezings of Ranunculus, epilithic, and plankton.
Pl. 11, Figs. 9 and 10.

*Cosmarium globosum Bulnheimi 1861.
(Actinotaenium subglobosum (Nordst.) Teiling 1954.)
West and West. 1908: 29, Pl. 68, Figs. 1 and 2.
L. 32-37 µm, W. 20-21 µm, Isth. 17-17.5 µm.
Common.
Sludge and benthic flocculum.
Pl. 7, Fig. 1.

*Cosmarium hammeri Reinsch 1867.
West and West. 1905: 181, Pl. 62, Figs. 20 and 21.
L. 30-33 µm, W. 23-26 µm, Th. 14-15 µm, Isth. 7.5-8 µm. The form was smaller than the Wests' which measured 40-50 µm x 27-35 µm.
Rare.
Sludge.
Pl. 7, Fig. 12.

*Cosmarium humile? (Gay) Nordst. var. striatum (Boldt) Schmidle 1895.
Taylor. 1934: 254, Pl. 51, Figs. 27-29.
West and West. 1908: 223, Pl. 85, Figs. 21 and 22.
L. 14 µm, W. 13 µm, Isth. 4-5 µm. Assignment to this species was questionable due to the lack of an undulate apex, but Taylor (1934) noted a reduction of this feature on his smaller forms.
Rare.
Benthic flocculum and tychoplankton.
Pl. 8, Fig. 6.

Cosmarium impressulum Elfving 1881. ∝
Krieger and Gerloff. 1965: 133, Pl. 29, Fig. 4.
L. 31 µm, W. 20 µm, Th. 14 µm, Isth. 8 µm.
Rare.
Squeezings of moss.

Pl. 7, Fig. 5.

Cosmarium impressulum Elfving 1881. β
Krieger and Gerloff. 1965: 133, Pl. 29, Fig. 4.
L. 17-20 µm, W. 13-15 µm, Th. 10 µm, Isth. 5 µm. The dimensions of this form and ∞ above were within the limits of the type. These, however, based on cell dimensions, formed two distinct groups without intermediates.
Rare.
Seepage and squeezings of Elodea.
Pl. 7, Fig. 7.

Cosmarium intermedium Delponte 1877.
West and West. 1908: 138, Pl. 76, Fig. 10.
L. 58-63 µm, W. 50-58 µm, Isth. 15-17 µm.
Rare.
Epilithic with Tetraspora lamellosa.
Pl. 10, Figs. 3 and 4.

*Cosmarium laeve Rabenhorst var. nov.?
West and West. 1908: 99, Pl. 73, Figs. 8-19.
L. 22 µm, W. 18 µm, Th. 9 µm, Isth. 6 µm. Cells were about 1 1/4 times as long as broad; semicells were semi-oblong, the apex was narrowly truncated and retuse as in the type, and sparsely punctate. The side view of the semicell was sub-ovate, and the apex was narrower than the base. The vertical view was elliptic. They differed from the type by the presence of a central pore.
Rare.
Benthic flocculum.
Pl. 7, Fig. 8.

*Cosmarium margaritatum (Lund.) Roy and Biss. forma subrotundata West and West 1912.
West and West. 1912: 19, Pl. 100, Fig. 1.
L. 70 µm, W. 56 µm, Th. 38 µm, Isth. 22 µm.
Rare.
Wet soil.
Pl. 10, Fig. 1.

*Cosmarium novae-semliae var. granulatum Schmidle 1898.
Croasdale. 1956: 42, Pl. 8, Fig. 9.
L. 18-19 µm, W. 17-19.5 µm, Th. 12 µm, Isth. 8 µm.
Rare.
Sludge and squeezings of moss.
Pl. 11, Fig. 5.

*Cosmarium ornatum var. perornatum Grönblad 1948.
Croasdale and Grönblad. 1964: 183, Pl. 13, Figs. 26 α27.
L. 35-40 µm, W. 35-38.5 µm, Th. 23 µm, Isth. 12-13 µm.
Rare.
Benthic flocculum and sludge.
Pl. 9, Fig. 1.

Cosmarium portianum Archer 1860.
West and West. 1908: 165, Pl. 80, Figs. 4-7.
L. 35-38 µm, W. 26-28 µm, Th. 20 µm, Isth. 10-12 µm.
Common.
Benthic flocculum and sludge.
Pl. 9, Fig. 5.

*Cosmarium pseudoarctoum Nordstedt 1879.
[Actinotaenium cruciferm (de Bary) Teiling 1954.]
West and West. 1908: 32, Pl. 118, Figs. 12-14, and Pl. 122, Figs. 40 and 41.
L. 22-23 µm, W. 15-16 µm, Isth. 13.5-14 µm.
Rare.
Sludge in Carex marsh.
Pl. 7, Fig. 6.

*Cosmarium pseudoprotuberans forma minus Kossinskaja 1936.
Croasdale. 1956: 47, Pl. 10, Fig. 5.
Hirano. 1968: 27, Pl. 5, Figs. 10-13 and 16.
L. 20 µm, W. 20.5 µm, Th. 10 µm, Isth. 7 µm.
Rare.
Plankton.
Pl. 8, Fig. 10.

*Cosmarium pseudopyramidatum Lund. var extensum (Nordst.) Gerloff 1965.
Krieger and Gerloff. 1965: 127, Pl. 26, Fig. 8.
L. 69-74 µm, W. 37-42 µm, Th. 25 µm, Isth. 15-16 µm. Most specimens closely fit the published description but several cells had a depressed apex as in var. excavatum.
Common.
Sludge and wet soil.
Pl. 8, Figs. 4 and 5.

*Cosmarium pseudoquadratulum Prescott and Scott 1952.
Prescott and Scott. 1952: 11, Fig. 5, No. 12.
L. 15-17.5 µm, W. 13-13.2 µm, Th. 7-8 µm, Isth. 3 µm.
Rare.
Sludge.
Pl. 7, Fig. 11.

Cosmarium pyramidatum Brébisson 1848.
Krieger and Gerloff. 1965: 121, Pl. 25, Fig. 3.
L. 70-74 µm, W. 38-48 µm, Th. 28 µm, Isth. 15-17 µm.
Cells had 1-3 pyrenoids per semicell.
Common.
Sludge and seepage.
Pl. 7, Figs. 3 and 4.

*Cosmarium quadratum Ralfs forma willei West and West 1908.
West and West. 1908: 59, Pl. 87, Figs. 21 and 22.
L. 52-61 µm, W. 28-33 µm, Isth. 16-20 µm.
Common.

Moss squeezings.
Pl. 8, Fig. 9.

*Cosmarium quinarium Lundell 1871.
West and West. 1908: 216, Pl. 85, Figs. 9 and 10.
L. 47-50 µm, W. 36-40 µm, Isth. 9-12 µm. Ours differed from the Wests' by having numerous small punctations about the central granules.
Common.
Sludge.
Pl. 9, Fig. 4.

*Cosmarium rectangulare Grun. var. hexagonium (Elfv.) West and West 1908.
West and West. 1908: 56, Pl. 80, Fig. 4.
L. 26-30 µm, W. 21-24 µm, Th. 14 µm, Isth. 7-9 µm.
Common.
Net tow in 15 centimeters of water, sludge, and net tow in Elodea.
Pl. 7, Fig. 2.

Cosmarium reniforme (Ralfs) Archer 1874.
West and West. 1908: 157, Pl. 79, Figs. 1 and 2, and Pl. 82, Fig. 15.
L. 45-49 µm, W. 40-42.5 µm, Th. 24 µm, Isth. 12-14 µm.
Common.
Squeezings of Elodea, squeezings of Ranunculus, and epilithic.
Pl. 9, Fig. 6.

*Cosmarium sexangulare Lundell 1871.
West and West. 1908: 81, Pl. 72, Fig. 3.
Prescott and Vinyard. 1965: 467, Pl. 11, Fig. 26.
L. 33-34 µm, W. 25-27 µm, Isth. 8 µm.
Rare.
Sludge and epilithic with Tetraspora lamellosa.
Pl. 7, Fig. 13.

*Cosmarium sexnotatum Gutw. var. tristriatum (Lütkem.) Schmidle 1895.
West and West. 1908: 228, Pl. 86, Figs. 8 and 9.
L. 17-18 µm, W. 17-19 µm, Th. 11 µm, Isth. 5-6 µm.
Common.
Benthic flocculum on gravel, sludge, and tychoplankton.
Pl. 11, Figs. 3 and 4.

*Cosmarium subarctoum (Lagerh.) Raciborski 1892.
Kreiger and Gerloff. 1962: 69, Pl. 16, Fig. 8.
L. 15 µm, W. 11-11.5 µm, Th. 7.5 µm, Isth. 8 µm.
Rare.
Sludge.
Pl. 7, Fig. 9.

Cosmarium subcrenatum Hantz. var. isthmochondrum Messikommer 1938.
Croasdale. 1973: 105, Pl. 16, Figs. 18-20.
L. 27.5-33 µm, W. 25-28.5 µm, Th. 17 µm, Isth. 7.5-10 µm.
Apex was smooth except for a few small scattered granules.
Common.

Plankton, tychoplankton, sludge, and benthic flocculum.
Pl. 9, Fig. 2, and Pl. 11, Figs. 1 and 2.

Cosmarium subspeciosum var. validius Nordst. 1887.
Croasdale and Grönblad. 1964: 187, Pl. 15, Figs. 1-3.
L. 72-79 µm, W. 48-59 µm, Th. 35-38 µm, Isth. 18-20 µm. These were quite similar to Croasdale and Grönblad's form.
Common.
Benthic flocculum and sludge.
Pl. 10, Figs. 8-11.

There apparently was a second form, larger than the first, lacking apical granules, and with 9 vertical series of granules on the central tumor.
L. 84-92 µm, W. 53-69 µm, Th. 42 µm, Isth. 20-22 µm.
Common.
Benthic flocculum on gravel shore.
Pl. 10, Figs. 6 and 7.

*Cosmarium subtumidum Nordst. var. minutum (Krieger) Krieger and Gerloff 1965.
Krieger and Gerloff. 1965: 164, Pl. 34, Fig. 4.
L. 16.5 µm, W. 14 µm, Th. 7.5 µm, Isth. 4 µm. Differed from typical variety in smaller L/W ratio. Considered a tropical species by Krieger and Gerloff (1965), who previously reported it in Java, Sumatra, and Brazil.
Very rare.
Epilithic with Tetraspora lamellosa.
Pl. 8, Fig. 13.

*Cosmarium tenue Arch. var. depressum Irénée-Marie 1952.
Irénée-Marie. 1952: 134, Pl. 12, Fig. 11.
Croasdale. 1956: 56, Pl. 5, Fig. 13.
L. 8-10 µm, W. 8-9 µm, Isth. 2.5-3 µm.
Very rare (probably often overlooked).
Plankton and sludge.
Pl. 8, Fig. 11.

*Cosmarium trachypleurum var. fallax Lütkemüller.
G. W. Prescott (personal correspondence).
L. 34 µm, W. 32 µm, Th. 21 µm, Isth. 4.5 µm.
Very rare.
Benthic flocculum.
Pl. 11, Fig.7.

*Cosmarium undulatum Corda ex Ralfs var. minutum Wittrock 1869.
Krieger and Gerloff. 1962: 41, Pl. 11, Fig. 11.
L. 29-34 µm, W. 21-25 µm, Th. 15-17 µm, Isth. 8-10 µm.
Rare.
Sqeezings of moss.
Pl. 8, Fig. 15.

*Cosmarium ungerianum (Näg.) de Bary 1858 forma.
G. W. Prescott (personal correspondence).

Croasdale. 1956: 59, Pl. 12, Figs. 9-14.
L. 55-60 µm, W. 41-47 µm, Th. 30-33 µm, Isth. 13-14 µm. Large, often low granules usually formed 4, sometimes 3 or 5 horizontal rows, each granule ringed with 6 distinct scrobiculations. Scrobiculations were circular, sometimes triangular on the central cell wall. Apex was scattered with scrobiculations or possibly large pores.
Very common.
Benthic flocculum and net tow in 15 centimeters of water.
Pl. 9, Figs. 7-9.

*Cosmarium venustum (Bréb.) Arch. var. excavatum W. West.
Grönblad. 1962: 478, Pl. 102, Fig. 13e.
L. 33-36 µm, W. 24-26 µm, Th. 16 µm, Isth. 7-10 µm.
Rare.
Sludge and wet soil.
Pl. 8, Fig. 3.

Desmidium Agardh

*Desmidium grevillii (Kütz.) de Bary.
Taylor. 1935: 219, Pl. 34, Fig. 11, and Pl. 49, Figs. 4 and 5.
L. 18-19 µm, W. 40-41 µm.
Very rare.
Squeezings of Elodea.
Pl. 5, Fig. 11.

Euastrum Ehrenberg

*Euastrum ansatum Ehrenb. var. pyxidatum Delponte 1876.
Kreiger. 1937: 489, Pl. 58, Fig. 7.
L. 54 µm, W. 27 µm, Isth. 9 µm.
Rare.
Benthic flocculum.
Pl. 13, Fig. 5.

*Euastrum ansatum Ehreb. var. triporum Krieger 1937.
Krieger. 1937: 492, Pl. 59, Fig. 8.
L. 84-90 µm, W. 42-44 µm, Wa. 20 µm, Th. 28 µm, Isth. 13 µm.
Rare.
Sludge.
Pl. 12, Fig. 3.

Euastrum bidentatum Näg. var. speciosum (Boldt) Schmidle 1898.
Krieger. 1937: 603, Pl. 85, Figs. 3-6.
L. 60-70 µm, W. 42-47 µm, Wa. 25 µm, Isth. 9-12 µm.
Common.
Sludge and net tow in 15 centimeters of water.
Pl. 12, Fig. 2.

*Euastrum denticulatum var. angusticeps Grönblad 1921.

Krieger. 1937: 584, Pl. 80, Figs. 18 and 19.
Prescott and Scott. 1945: 236, Pl. 2, Fig. 13.
L. 21.5-23 µm, W. 15.5-18 µm, Wa. 13 µm, Th. 13 µm, Isth. 5 µm. Granulation best fit Prescott and Scott's species description, but the lateral view was of the variety.
Rare.
Sludge and benthic flocculum.
Pl. 12, Fig. 5.

*Euastrum didelta Ralfs 1844.
Krieger. 1937: 517, Pl. 67, Figs. 1-3.
L. 115-128 µm, W. 65-70 µm, Wa. 28-29 µm, Isth. 18 µm.
Rare.
Benthic flocculum on gravel shore.
Pl. 13, Fig. 6.

Euastrum elegans (Bréb.) Kützing 1845.
Krieger. 1937: 591, Pl. 81, Figs. 14-18.
L. 30-33 µm, W. 21-24 µm, Wa. 15 µm, Isth. 4-7 µm.
Common.
Benthic flocculum, squeezings of Ranunculus, and net tow in 15 centimeters of water.
Pl. 13, Fig. 4.

Euastrum gemmatum Bréb. var. taftii Prescott (1977).
Prescott, Croasdale and Vinyard. 1977: 62, Pl. 76, Fig. 22, and Pl. 82, Figs. 3, 5.
L. 48-57 µm, W. 37-40 µm, Isth. 10-11.5 µm.
Rare.
Sludge.
Pl. 13, Figs. 7 and 8.

*Euastrum obesum Joshua 1886.
Krieger. 1937: 495, Pl. 59, Figs. 9 and 10.
L. 51-58 µm, W. 30-34 µm, Wa. 17-18 µm, Th. 17 µm, Isth. 11-12 µm. They were densely and delicately punctate.
Rare.
Sludge.
Pl. 12, Fig. 4.

Euastrum oblongum (Grev.) Ralfs 1844.
Krieger. 1937: 526, Pl. 70, Figs. 3-6.
L. 136-186 µm, W. 68-92 µm, Wa. 40 µm, Th. 42 µm, Isth. 19-28 µm.
Common.
Benthic flocculum on gravel shore, sludge, and wet soil.
Pl. 12, Fig. 1.

*Euastrum verrucosum Ehrenb. var. perforatum Grönblad 1920.
Krieger. 1937: 649, Pl. 95, Fig. 6.
L. 118 µm, W. 106 µm, Wa. 36 µm, Isth. 25 µm. They lacked upper lateral lobes.
Rare.

Benthic flocculum on gravel shore.
Pl. 13, Fig. 1.

Euastrum verrocosum Ehrenb. var. rhomboideum Lundell 1871.
Krieger. 1937: 650, Pl. 96, Fig. 2.
L. 100-120 µm, W. 86-94 µm, Wa. 32 µm, Th. 51 µm, Isth. 20-22 µm . They had pores on the central protuberance.
Common.
Net tow in 15 centimeters of water and benthic flocculum on gravel shore.
Pl. 13, Fig. 2.

*Euastrum verrucosum Ehrenb. var. vallesiacum Viret 1909.
Krieger. 1937: 653, Pl. 96, Fig. 8.
L. 83-90 µm, W. 68-73 µm, Isth. 20-21 µm. Interlobular surfaces were smooth.
Rare.
Seepage.
Pl. 13, Fig. 3.
All three varieties of E. verrucosum were found in the same or similar habitats within Rae Lake 2. Perhaps the validity of these varieties should be re-examined.

Micrasterias Agardh

Micrasterias americana (Ehrenb.) Ralfs 1848.
West and West. 1905: 116, Pl. 53, Figs. 1-3.
L. 168-170 µm, W. 150-153 µm, Wa. 76-78 µm, Isth. 26-30 µm. They were larger than the Wests' in all dimensions.
Rare.
Benthic flocculum and benthic flocculum on gravel shore.
Pl. 12, Fig. 8.

*Micrasterias conferta Lundell 1871.
West and West. 1905: 88, Pl. 43, Figs. 4-8.
L. 96-110 µm, W. 77-85 µm, Isth. 21-25 µm. Some cells with apparent concretions of mucilage, affected both semicells.
Rare.
Sludge.
Pl. 14, Figs. 5 and 6.

*Micrasterias denticulata var. angulosa (Hantzs.) West and West 1902.
West and West. 1905: 107, Pl. 2, Figs. 3 and 4.
L. 262-277 µm, W. 200-214 µm, Th. 60 µm, Isth. 35-37.5 µm. Walls were smooth and non-punctate.
Rare.
Benthic flocculum and sludge.
Pl. 14, Figs. 1 and 8.

*Micrasterias muricata var. tumida West and West 1896.
Krieger. 1937: 75, Pl. 119, Fig. 7.

L. 172 µm, W. 120 µm, Wa. 80-90 µm, Th. 45 µm , Isth. 30 µm. Walls were punctate and the tips of each process had five points. The isthmus was almost twice the width of Krieger's.

Very rare.

Benthic flocculum.

Pl. 14, Fig. 7.

Micrasterias pinnatifida (Kütz.) Ralfs 1848.

West and West. 1905: 80, Pl. 41, Figs. 7-11 and 13.

L. 62-73 µm, W. 58-74 µm, Wa. 39-50 µm, Isth. 15 µm. Lobular attenuations were bluntly granular or sharply extended.

Common.

Seepage and sludge.

Pl. 14, Figs 2 and 3.

*Micrasterias rotata (Grev.) Ralfs 1844.

West and West. 1905: 102, Pl. 48, Figs. 1-6.

L. 250-300 µm, W. 225-273 µm, Isth. 37-40 µm.

Common.

Benthic flocculum and squeezings of Elodea.

Pl. 14, Fig. 4.

*Micrasterias rotata forma evoluta Turner 1893.

West and West. 1905: 104, (not illustrated).

L. 266-292 µm, W. 235-295 µm, Isth. 38-42 µm. They had a tooth on one side of the median emargination and a swelling on the other instead of a tooth on both sides. The tooth on the front side was opposite a swelling on the back side.

Common.

Benthic flocculum on gravel shore and squeezings of Elodea.

Pl. 12, Fig. 6.

Micrasterias truncata var. neodamensis (Braun) Dick 1926.

Prescott and Scott. 1952: 250, Pl. 7, Fig. 5.

L. 110 µm, W. 95 µm, Isth. 23 µm. They were punctate and almost twice the size of Prescott and Scott's (L. 66 µm, W. 65 µm, Isth. 16 µm).

Rare.

Seepage.

Pl. 12, Fig. 7.

Penium Brebisson

*Penium spirostriolatum Barker 1869.

West and West. 1904: 88, Pl. 9, Figs. 1-8.

L. 185 µm, W. (at constriction) 25 µm, Wa. 16 µm.

Very rare.

Sludge.

Pl. 5, Figs. 8 and 9.

Pleurotaenium Nägeli

Pleurotaenium ehrenbergii (Bréb.) de Bary 1858.
Kreiger. 1937: 410, Pl. 42, Figs. 4-8.
L. 515-650 µm, W. (at constriction) 26-47 µm, Wa. 22-27 µm.
Very common.
Benthic flocculum on gravel and squeezings of Ranunculus.
Pl. 5, Figs. 2 and 3.

*Pleurotaenium trabecula var. elongatum Cedergren 1913.
Krieger. 1937: 399, Pl. 40, Fig. 5.
L. 620-700 µm, W. (at constriction) 25-35 µm, Wa. 20-28 µm.
Rare.
Plankton, net tow in Elodea, and epilithic with Nostoc parmelioides.
Pl. 5, Fig. 4.

*Pleurotaenium trabecula (Ehrenb.) Naeg. var. maximum (Reinsch) Roll 1927.
Krieger. 1937: 400, Pl. 40, Fig. 8.
L. 670-800 µm, W. (at constriction) 50-52 µm, Wa. 38-45 µm. Margins were undulate with large basal inflation (70 µm in diameter). Two plants were found; one which may be abnormal had a swollen apex and large inflation (80 µm) above the basal inflation.
Very rare.
Benthic flocculum on gravel.
Pl. 5, Fig. 1.

Spondylosium Brébisson

*Spondylosium planum (Wolle) West and West 1912.
West, West and Carter. 1923: 222, Pl. 160, Figs. 23-25.
L. 8-10 µm, W. 10-13 µm, Th. 5 µm, Isth. 4-6 µm.
Rare.
Sludge in Carex marsh, squeezings of Ranunculus, and net tow in 15 centimeters of water.
Pl. 18, Fig. 7.

*Spondylosium pulchellum Archer 1858.
West, West and Carter. 1923: 227, Pl. 161, Figs. 1-3.
L. 11-13 µm, W. 9-11 µm, Wa. 7-7.5 µm, Isth. 3.5-4.5 µm.
Very rare.
Sludge.
Pl. 18, Fig. 6.

Staurastrum Meyen

Many species of Staurastrum were transferred to the genus Staurodesmus by Teiling (1967). Since his nomenclature has been accepted by many students of the desmids, both names, wherever applicable, are given in this text.

*Staurastrum acarides Nordstedt 1872.
West, West and Carter. 1923: 73, Pl. 140, Figs. 6 and 7.
L. 40 µm, W. 30 µm, Wa. 20 µm, Isth. 15 µm.

Very rare.
Wet soil.
Pl. 16, Fig. 4.

Staurastrum alternans Brébisson 1848.
West and West. 1912: 170, Pl. 126, Figs. 8 and 9.
L. 28-32 µm, W. 32-35 µm, Isth. 10-13 µm. Granules were sharp and distinct.
Common.
Sludge and benthic flocculum on gravel.
Pl. 17, Figs. 9 and 11.

*Staurastrum anatinum Cooke and Wills 1880.
West, West and Carter. 1923: 142, Pl. 147, Fig. 1.
L. 43-50 µm, W. 85-100 µm, Isth. 12 µm. They were similar to the smaller form described by the Wests and Carter.
Common.
Squeezings of Ranunculus, plankton, net tow in creek, and net tow in 15 centimeters of water.
Pl. 16, Figs. 1 and 2.

Staurastrum arctiscon (Ehrenb.) Lundell 1871.
West, West and Carter. 1923: 193, Pl. 157, Fig. 5.
L. (body) 63-66 µm, L. (+ processes) 122-124 µm, W. (body) 43-45 µm, W. (+ processes) 108-130 µm, Isth. 23 µm.
Very common.
Occurred in most habitats.
Pl. 18, Fig. 1.

Staurastrum brebissonii Archer 1861.
West, West and Carter. 1923: 61, Pl. 137, Figs. 4 and 5.
L. (body) 43 µm, W. (body) 47-56 µm, Spines 4-5 µm, Isth. 15 µm.
Rare.
Benthic flocculum on gravel.
Pl. 15, Fig. 13.

*Staurastrum breviaculeatum Smith 1924.
Smith. 1924: 78, Pl. 70, Figs. 10-18.
L. (body) 43-44 µm, W. (body) 38-40 µm, Spines 3-7 µm, Isth. 13-16 µm. Length and number of spines were variable in our material.
Very common.
Sludge and benthic flocculum.
Pl. 17, Figs. 1-3.

Staurastrum brevispinum Brébisson 1848.
(Staurodesmus brevispina (Bréb.) Croasdale 1957.)
Smith. 1924: 68, Fig. 7, A-E.
L. 43 µm, W. 42 µm, Isth. 13 µm.
Rare.
Squeezings of Elodea and plankton.
Pl. 18, Fig. 3.

*Staurastrum crenulatum (Näg.) Delponte 1887.

West, West and Carter. 1923: 110, Pl. 143, Figs. 9-13.
L. 23 µm, W. 37 µm, Isth. 5 µm.
Very rare.
Squeezings of Ranunculus.
Pl. 17, Fig. 10.

Staurastrum cuspidatum Brébisson 1840.
[Staurodesmus cuspidatus (Bréb.) Teiling 1967.]
Smith. 1924: 74, Pl. 68, Figs. 27-34.
L. 22 µm, W. (- spines) 19-20 µm, W. (+ spines) 38-40 µm, Isth. 5 µm, Spines 9-11 µm.
Common.
Benthic flocculum.
Pl. 17, Fig. 8.

Staurastrum dickiei Ralfe var. circulare Turner 1893.
[Staurodesmus convergens var. laportei Teiling 1967.]
West, West and Carter. 1923: 5, Pl. 124, Fig. 16.
L. 30 µm, W. 33 µm, Isth. 11 µm.
Very rare.
Seepage into Lake 2.
Pl. 15, Fig. 4.

*Staurastrum grande Bulnh. var. angulosum Grönblad 1920.
[Staurodesmus grandis (Bulnh.) Teiling 1967.]
Grönblad. 1920: 66, Pl. 3, Figs. 107 and 108.
L. 44-45 µm, W. 40-43 µm, Isth. 13-19 µm. They were smaller than Grönblad's and the cell wall was thickened at the basal angles.
Common.
Sludge.
Pl. 15, Fig. 6.

Staurastrum kurilense Okada 1924.
Vinyard. 1951: 44, Pl. 11, Figs. 3 and 4.
Carter. 1935: 171, Figs. 12 and 13.
L. 36 µm, W. 32 x 60 µm, Isth. 12 µm. They were sparsely punctate. Carter reported this plant as Staurastrum natator West var. rhomboideum. Vinyard noted that Okada's description had priority.
Rare.
Sludge and benthic flocculum.
Pl. 17, Figs. 6 and 7.

Staurastrum kurilense fa. triquetra Carter 1935.
Vinyard. 1951: 45, Pl. 11, Fig. 5.
Carter. 1935: 171, Figs. 33 and 34.
L. 46 µm, W. 43-46 µm, Isth. 14 µm. Vinyard (1951) noted that, "this is a 3-angled form of the species figured by Wailes (1930b), and described by Carter as a form of St. natator". See St. kurilense above. Mine differed in the number and position of the apical verrucose processes, with 9 instead of 6, 1 above each short lateral process and 2 above each long lateral process. Cell wall was sparsely punctate.
Very rare.

Benthic flocculum.
Pl. 17, Figs. 4 and 5.

Staurastrum margaritaceum (Ehrenb.) Meneghini 1840.
West, West and Carter. 1923: 131, Pl. 150, Figs. 5-9.
L. 33 µm, W. 29-34 µm, Isth. 10 µm.
Very rare.
Benthic flocculum on gravel shore.
Pl. 15, Fig. 14.

*Staurastrum orbiculare Ralfs 1845.
West and West. 1912: 155, Pl. 124, Figs. 10 and 11.
L. 45-48 µm, W. 28-37 µm, Isth. 11-12 µm.
Rare.
Sludge and benthic flocculum.
Pl. 15, Fig. 9.

*Staurastrum pachyrhynchum Nordstedt 1875.
[Staurodesmus pachyrhynchus (Nordst.) Teiling 1963.]
West and West. 1912: 151, Pl. 121, Figs. 8 and 9.
L. 33-35 µm, W. 27-30 µm, Isth. 10-11 µm.
Common.
Sludge.
Pl. 15, Fig. 7.

*Staurastrum polonicum Raciborski 1884.
G. W. Prescott (personal correspondence).
L. 40-43 µm, W. 28-32 µm, Isth. 18-20 µm. Plants were 7 or 8 angled.
Common.
Squeezings of moss and wet soil.
Pl. 15, Fig. 10.

*Staurastrum polymorphum Brébisson 1848.
West, West and Carter. 1923: 125, Pl. 142, Fig. 24, and Pl. 143, Figs. 1-3.
L. 20-22 µm, W. 27-28 µm, Isth. 7.5 µm.
Rare.
Sludge and epilithic.
Pl. 15, Fig. 12.

*Staurastrum punctulatum Brébisson 1848.
Prescott and Scott. 1951: 64, Fig. 14, No. 18.
West and West. 1912: 179, Pl. 127, Figs. 8-11, 13, and 14.
L. 30-34 µm, W. 32-40 µm, Isth. 10-12 µm. Granules were flattened.
Common.
Benthic flocculum on gravel shore and sludge in Elodea.
Pl. 15, Fig. 11.

*Staurastrum punctulatum Breb. var. kjellmani Wille 1879.
West and West. 1912: 182, Pl. 127, Figs. 13, 17-19, 21, and 22.
L. 32.5-43 µm, W. 29-38 µm, Isth. 12-17.5 µm. The semicells were 3 and 4-angled and the 4-angled forms were larger.

Common.
Epilithic, benthic flocculum, and benthic flocculum on gravel shore.
Pl. 16, Figs. 3, 5, and 6.

*Staurastrum pyramidatum West and West forma nov.?
G. W. Prescott (personal correspondence).
L. 43-50 µm, W. 46-47 µm, Isth. 11-12 µm. Differed from the type in smaller dimensions; angles were more sharply rounded, slightly incurved, and smooth tipped. Granules were short and blunt at base and angles of semicell, becoming longer to form sharp conical spines towards the apex.
Rare.
Seepage and sludge.
Pl. 15, Fig. 8.

Staurastrum sebaldii var. ornatum Nordstedt 1873.
West, West and Carter. 1923: 167, Pl. 148, Fig. 7.
Sieminska. 1965: 117, Pl. 7, Figs. 18-20.
L. 80-90 µm, W. 105-150 µm, Isth. 23-24 µm. Three and 4-angled forms were present.
Common.
Net tow in 15 centimeters of water.
Pl. 16, Figs. 7, 8, and 10.

*Staurastrum spongiosum Brebisson 1848.
West, West and Carter. 1923: 76, Pl. 140, Fig. 14.
L. 52 µm, W. 45-48 µm, Isth. 13-16 µm.
Very rare.
Sludge and squeezings of moss.
Pl. 15, Figs. 1-3.

*Staurastrum subavicula W. and G. S. West 1894.
West, West and Carter. 1923: 181, Pl. 155, Fig. 10.
L. about 34 µm(- spines), L. about 42 µm(+ spines), W. 33 µm, Isth. 12 µm. Differed from the type in 4 series of granules on angles instead of 2, angles tipped with 3-4 spines instead of 2, and center of apex was punctate.
Very rare.
Sludge.
Pl. 15, Fig. 5.

Staurastrum vestitum Ralfs 1848.
West, West and Carter. 1923: 258, Pl. 151, Figs. 9-11, and Pl. 152, Figs. 5 and 6.
L. 35-42 µm, W. 56-60 µm, Isth. 11-13 µm.
Common.
Benthic flocculum.
Pl. 16, Fig. 9.

Tetmemorus Ralfs

*Tetmemorus Laevis (Kütz.) Ralfs 1848.
West and West. 1904: 222, Pl. 32, Figs. 11-16.

L. 83-84 µm, W. 26 µm, Isth. 23 µm.
Very rare.
Wet soil.
Pl. 6, Fig. 17.

Xanthidium Ehrenber

*Xanthidium subhastiferum West 1892.
West and West. 1912: 56, Pl. 106, Figs. 5-9.
L. (- spines) 50-56 µm, L. (+ spines) 60-66 µm, W. (- spines) 50-52 µm, W. (+ spines) 81-96 µm, Th. 28-30 µm, Isth. 11.5-14.5 µm. Cells with 3 pairs of spines per semicell reached a maximum length of 96 µm.
Very common.
Plankton, benthic flocculum, tychoplankton, and sludge.
Pl. 18, Fig. 2.

Charales

Characeae

Nitella Agardh

Nitella sp.
Mature reproductive structures were lacking. This plant was introduced in 1919 for the purpose of establishing a rainbow trout fishery (Coleman 1925).
Benthic, fringing beds of Elodea nuttalii at depths less than 1 meter.

EUGLENOPHYTA

Euglenales

Euglenaceae

Euglena Ehrenberg

Euglena spirogyra var. marchica Lemmerman.
Johnson. 1944: 112, Fig. 11-D.
Cells 150 µm long, 23 µm wide, spine 15 µm long. Determination based on external characteristics only.
Very rare.
Sludge.
Pl. 4, Fig. 4.

Lepocinclis Perty

Prescott. 1962: 406, Pl. 89, Figs. 7 and 15.
Cells 60 µm long, 39 µm in diameter. They had two large paramylon rings and unlike Prescott's, had typical spiral striations.
Very rare.
Epilithic with Tetraspora lamellosa.
Pl. 4, Fig. 6.

Lepocinclis fusiformis (Carter) Lemmermann 1901.
Prescott. 1962: 406, Pl. 89, Figs. 1-4.
Cells 31 µm long, 22 µm in diameter with 2 U-shaped paramylon bodies on opposite sides of the cell, broadly ovate, without a caudus. Flagellum extended through a truncated apical protrusion. Periplast had striations gently spiralling to the left.
Rare.
Sludge.
Pl. 4, Fig. 14.

*Lepocinclis fusiformis var. major Fritisch and Rich 1930.

Phacus Dujardin

Phacus sp.
Cells 26 µm long, 20 µm wide, broadly ovoid, with a short caudus angling slightly to the right (as seen in ventral view). Caudus 4 µm long. Periplast had very fine spiral striations. Paramylon was in the form of 10 rings. Flagellum was 1 1/2 times as long as the cell.
Very rare.
Sludge.
Pl. 4, Fig. 11.

Trachelomonas Ehrenberg

Trachelomonas bacillifera Playf. var. minima Playfair.
Huber-Pestalozzi. 1955: 303, Fig. 556.
Test about 23 µm long, 16-19 µm in diameter. Test broadly oval and brown. Flagellum aperture had a very short collar. Wall was densely covered with stout, blunt spines.
Rare.

*Trachelomonas horrida Palmer 1905.
Prescott. 1962: 415, Pl. 84, Fig. 1.
Test 40 µm long, 25 µm in diameter.
Very rare.
Sludge.
Pl. 4, Fig. 10.

*Trachelomonas lacustris Drezepolski 1925.
Prescott. 1962: 415, Pl. 83, Figs. 14 and 15, and Pl. 85, Fig. 15.
Test 27 µm long, 14 µm in diameter.
Very rare.
Sludge.
Pl. 4, Fig. 16.

Sludge.
Pl. 4, Fig. 17.

PYRROPHYTA

Dinokontae

Peridinium Ehrenberg

Peridinium willei Huitfeld-Kaas 1900.
Prescott. 1962: 434, Pl. 91, Figs. 22-25.
Cells 50-60 µm long, 52-60 µm wide, 40-45 µm thick.
Very common.
Plankton and tychoplankton.
Pl. 4, Fig. 15.

CHRYSOPHYTA

Mischococcales

Chlorobotrydaceae

Ducellieria Teiling

**Ducellieria chodatii* (Ducell.) Teiling.
Bourrelly. 1968: 197, Pl. 39, Fig. 7, and Pl. 40, Fig. 1.
Cells 12-16 µm long (- spine), 11 µm in diameter, spine 5-6 µm long. Colonies of 9 and 16 cells were observed, 45 µm and 68 µm in diameter respectively.
Very rare.
Benthic flocculum.
Pl. 5, Figs. 10 and 15.

Ochromonadales

Dinobryaceae

Dinobryon Ehrenberg

**Dinobryon cylindricum* Imhof 1883.
Prescott. 1962: 378, Pl. 107, Fig. 1.
Lorica 40-50 µm long, mouth 7-12 µm in diameter.

Rare.
Plankton and tychoplanton.
Pl. 5, Fig. 12.

CYANOPHYTA

Chroococcales

Chroococcaceae

Aphanocapsa Nägeli

*Aphanocapsa elachista var. conferta West and West 1912.
Prescott. 1962: 453, Pl. 101, Figs. 10 and 11.
Cells 1.5-2 µm in diameter. Colonies up to 110 µm in diameter.
Common.
Tychoplankton and benthic flocculum.
Pl. 19, Fig. 2.

*Aphanocapsa pulchra (Kütz.) Rabenhorst 1865.
Prescott. 1962: 454, Pl. 101, Fig. 14.
Cells 3-4 µm in diameter.
Rare.
Plankton in seepage.
Pl. 19, Fig. 3.

Aphanothece Nägeli

Aphanothece stagnina (Spreng.) Braun 1865.
Desikachary. 1959: 137, Pl. 21, Fig. 10.
Prescott. 1962: 469, Pl. 103, Figs. 14-16.
Cells 5-6.5 µm long, 3-3.5 µm in diameter. Colonies microscopic.
Common.
Benthic flocculum and sludge.
Pl. 19, Fig. 1.

Chroococcus Nägeli

*Chroococcus prescottii Drouet and Daily 1942.
Prescott. 1962: 450, Pl. 100, Fig. 13.
Cells 5-8 µm in diameter. Colonies were of 2, 4, 8, rarely 16 and 32 cells. Larger colonies were in loose groups of 8 cells.
Common.
Tychoplankton, sludge, and benthic flocculum.
Pl. 19, Fig. 4.

Chroococcus turgidus (Kütz.) Nägeli 1849.

Prescott. 1962: 450, Pl. 100, Fig. 19.
Cells hemispherical, 8-30 µm in diameter.
Common.
Plankton, tychoplankton, sludge, and benthic flocculum.
Pl. 19, Fig. 5.

Coelosphaerium Nägeli

Coelosphaerium kutzingianum Nägeli 1849.
Prescott. 1962: 470, Pl. 106, Fig. 2.
Cells 2-3.5 µm in diameter.
Rare.
Net tow in 15 centimeters of water.
Pl. 19, Fig. 7.

*Coelosphaerium pallidum? Lemmermann 1898.
Prescott. 1962: 471, Pl. 106, Fig. 3.
Cells 1.5-3 µm in diameter, 3.5-5 µm long. The cells were larger than Prescott's, within the range of C. naegelianum Unger but without pseudovacules and radiating fibrillar concretions. Colonies were spherical, ovate or lobed.
Common.
Benthic flocculum, plankton, and tychoplankton.
Pl. 19, Fig. 6.

Dactylococcopsis Hansgirg

*Dactylococcopsis smithii Chodat and Chodat 1925.
Prescott. 1962: 465, Pl. 105, Figs. 3 and 4.
Cells 8-10 µm long, 2-2.4 µm in diameter, smaller than the type but properly proportioned
Very rare.
Sludge.
Pl. 19, Fig. 10.

Eucapsis Clements and Shantz

*Eucapsis alpina Clements and Schantz var. minor Skuja.
Sieminska. 1965: 99, Pl. 1, Fig. 1.
Cells 2-3 µm in diameter, usually 32 cells in a distinct colony 8 µm x 12 µm x 15 µm, often with several colonies adjoined.
Rare.
Benthic flocculum.
Pl. 19, Fig. 13.

Gloeothece Nägeli

*Gloeothece rupestris? (Lyngb.) Bornet 1880.

Prescott. 1962: 462, Pl. 103, Figs. 2 and 3.
Cells up to 8 µm long, 2.5-3 µm in diameter. The cells were about half the size of Prescott's, otherwise they followed his description closely.
Plankton.
Pl. 19, Fig. 9.

Merismopedia Meyen

Merismopedia elegans A. Braun 1849.
Prescott. 1962: 459, Pl. 101, Fig. 1.
Cells 7-8 µm long, 4-6 µm in diameter. Colonies seldom had more than 100 cells.
Very common.
Sludge and benthic flocculum.
Pl. 19, Fig. 11.

Microcystis Kützing

Microcystis aeruginosa Kützing 1846.
Prescott. 1962: 456, Pl. 102, Figs. 1-4.
Cells (2.5)-3-4 µm in diameter. Colonies were clathrate except when very small. They lacked pseudovacules, a character usually noted for the species when reported as a euplankter. The colonies were always collected in sludge and benthic flocculum with the exception of one net tow in moving water where the colonies may have been carried into the plankton.
Common.
Benthic flocculum, tychoplankton, squeezings of Elodea and Ranunculus, and sludge.
Pl. 19, Figs. 8 and 12.

Oscillatoriales

Oscillatoriaceae

Lyngbya Agardh

Lyngbya aerugineo-caerulea (Kütz.) Gomont 1892.
Prescott. 1962: 498, Pl. 111, Figs. 10 and 11.
Cells 5 µm in diameter, 2.5-3 µm long. Sheath 6 µm in diameter. Filaments were solitary.
Rare.
Benthic flocculum, plankton, and seepage.
Pl. 20, Fig. 4.

Oscillatoria Vaucher

*Oscillatoria agardhii Gomont 1892.

Desikachary. 1959: 235, (not illustrated).
Prescott. 1962: 484, Pl. 108, Figs. 15 and 16.
Cells 6.5-7 µm in diameter, 2.5-7 µm long. Apical cell was rounded, without a calyptra. Trichomes were scattered.
Common.
Plankton, tychoplankton, sludge, and benthic flocculum.
Pl. 20, Fig. 2.

*Oscillatoria limnetica Lemmermann 1900.
Prescott. 1962: 488, Pl. 109, Fig. 16.
Cells 1.5-2 µm in diameter, 3-5 µm long.
Common.
Tychoplankton.
Pl. 20, Fig. 3.

*Oscillatoria tenuis C. A. Agradh 1813,
Prescott. 1962: 491, Pl. 110, Figs. 8, 9, and 14.
Cells 4-4.5 µm in diameter, 2.5-3 µm long, slightly constricted at the crosswalls. Trichomes were loosely intermingled or scattered.
Common.
Tychoplankton.
Pl. 20, Fig. 5.

*Oscillatoria tenuis var. natans Gomont 1892.
Prescott. 1962: 491, Pl. 110, Figs. 10 and 11.
Cells 9-11 µm in diameter, 2.5-5 µm long, slightly constricted at crosswalls. Trichomes were scattered.
Common.
Tychoplankton and benthic flocculum.
Pl. 20, Fig. 1.

Phormidium Kützing

*Phormidium corium (Agardh) Gomont 1890.
Prescott. 1962: 494, (not illustrated).
Tilden. 1910: 101, Pl. 4, Figs. 71 and 72.
Cells 2.5-6 µm long, 4 µm in diameter. Filaments were parallel, forming compact bundles with ends free bending away from the bundle.
Rare.
Sludge.
Pl. 20, Fig. 6.

Nostocales

Nostocaceae

Anabaena Bory

*Anabaena oscillariodes Bory 1822.
Desikachary. 1959: 417, Pl. 71, Fig. 7.
Cells 4-5 µm long, 3.5-6 µm in diameter. Heterocysts spherical, 6.5-8

µm in diameter. Akinetes 20-37 µm long, 9 µm in diameter, mostly on just one side of the heterocyst.

Sometimes the filament had a watery sheath 14 µm in diameter.

Very common.

Sludge, wet soil, plankton, and benthic flocculum.

Pl. 20, Fig. 13.

*Anabaena sphaerica Bornet and Flahault 1888.

Desikachary. 1959: 393, (not illustrated).

Cells 6.5-8 µm in diameter, spherical, shorter, or longer than wide. Heterocysts spherical, 7.5-8.5 µm in diameter. Akinetes on one or both sides of heterocyst, 8.5-9 µm in diameter, 10 µm long. Filaments were scatterd.

Common.

Seepage and benthic flocculum.

Pl. 20, Fig. 9.

Cylindrospermum Kützing

*Cylindrospermum alatosporum Fritsch 1918.

Desikachary. 1959: 362, Pl. 64, Fig. 9.

Cells 3 µm in diameter, 3.5-7.5 µm long. Heterocysts 3.5-5.5 µm in diameter, 5.5-8 µm long. Akinetes 9-10 µm in diameter, 17-19 µm long, walls thick and punctate with age. Filaments were usually solitary. The cells, like the Indian form, were narrower than the type. Akinete lacked a thick yellowish inner wall.

Rare.

Sludge.

Pl. 20, Fig. 10.

Nostoc Vaucher

*Nostoc parmelioides Kützing 1843.

Desikachary. 1959: 389, Pl. 70, Fig. 3.

Cells 4-7 µm long, 3-5 µm in diameter. Heterocysts 7.5-8.5 µm long, 6.5-8.5 µm in diameter. Thallus lacked central radiating filaments. The arrangement of cells into trichomes varied within the same thallus; cells at periphery of thallus were adjacent to one another forming continuous trichomes with yellow sheaths, interior cells were 1-2 cell diameters apart, forming loose trichomes with very wide sheaths visible only after staining. Colonies were irregularly lobed, macroscopic, forming epilithic expanses.

Common.

Epilithic.

Pl. 20, Figs. 14-16 and 18.

*Nostoc paludosum Kützing 1850.

Desikachary. 1959: 375, Pl. 69, Fig. 2.

Cells 3-5 µm in diameter, 3.5-5 µm long. Heterocysts spherical to slightly ovate, 5-7 µm in diameter. Akinetes ovate 4-5 µm in diameter, 5.5-8 µm long.

Very common.

Plankton, tychoplankton, benthic flocculum, seepage, and sludge.
Pl. 20, Fig. 20.

Scytonemataceae

Scytonema Agardh

*Scytonema mirabile (Dillw.) Bornet 1889.
Prescott. 1962: 535, Pl. 124, Figs. 7 and 8.
Cells 5-16 µm long, 4-6 µm in diameter. Filaments 12-19 µm in diameter. Heterocysts 9-12 µm long, 6-10 µm in diameter. They seldom formed double false branches, diverging lamellations were found rarely and only in older sheaths.
Common.
Tychoplankton, sludge, and benthic flocculum.
Pl. 20, Fig. 19.

Tolypothrix Kützing

*Tolypothrix distorta Kützing 1843.
Prescott. 1962: 537, Pl. 125, Figs. 5 and 6.
Cells 2.5-7 µm long, 7-10 µm in diameter. Sheath sometimes lamellate, 10-17 µm in diameter, old sheaths were as wide as 25 µm in diameter. Heterocysts spherical, to ovate, 9-20 µm long, 9-11 µm in diameter. Presence of a lamellate sheath was inconsistent with the type, otherwise, they agreed.
Common.
Plankton, sludge, tychoplankton, and benthic flocculum.
Pl. 20, Figs. 7 and 8.

Rivulariaceae

Calothrix Agardh.

*Calothrix epiphytica West and West 1897.
Prescott. 1962: 553, Pl. 132, Figs. 2 and 3.
Desikachary. 1959: 543, (not illustrated).
Basal cells 3-5 µm in diameter, shorter than wide. Filaments 5-6.5-(9) µm in diameter. Sheath sometimes extended the entire length of the trichome. Heterocysts basal, 3-5 µm in diameter.
Common.
Epiphytic on Rhizoclonium crassipellitum.
Pl. 20, Figs. 11 and 12.

*Calothrix fusca (Kütz.) Bornet and Flahault 1886.

Prescott. 1962: 553, Pl. 132, Figs. 4 and 5.

Basal cells 9-10 µm in diameter, length 1 3 the width. Sheath 11-16 µm wide. Heterocyst basal, hemispherical, 7-7.5 µm in diameter, slightly smaller than Prescott's.

Rare.

Associated with the mucilage of Nostoc parmelioides and Tetraspora lamellosa.

Pl. 20, Fig. 17.

APPENDIX B

SPECIES ILLUSTRATIONS

The line drawn beside each illustration is equal to 10 micrometers.

PLATE 1

Figure		Page
1	Eudorina elegans	25
2	Chlamydomonas angulosa?	25
3	Pandorina morum	25
4	Tetraspora lamellosa	25
5	Asterococcus limneticus	26
6	Tetraëdron gracile	26
7	Gloeocystis vesiculosa	26
8	Ankistrodesmus gelifactus	27
9	Ankistrodesmus falcatum	27
10	Gloeocystis gigas	26
11	Gloeocystis ampla	26
12	Sphaerocystis schroeteri	27

PLATE 1



PLATE 2

Figure		Page
1	*Eremosphaera viridis*	27
2	*Oocystis arctica*	28
3	*Oocystis parva*	28
4	*Oocystis borgei*	28
5,6	*Oocystis elliptica* ?	28
7	*Oocystis pusilla* ?	28
8	*Oocystis* sp.	28
9	*Golenkinia paucispina*	29
10	*Kirchneriella lunaris* v. *irregularis*	27
11	*Dictyosphaerium pulchellum*	29
12	*Quadrigula lacustris*	29
13	*Botryococcus braunii*	29

PLATE 2



PLATE 3

Figure		Page
1	Coelastrum cambricum v. intermedium	30
2,3	C. microporum	30
4	C. printzii	30
5	Crucigenia quadrata	30
6	C. rectangularis	30
7	Coelastrum proboscideum	30
8	Enallax alpina	31
9	Scenedesmus bijuga v. alternans	31
10	S. quadricauda	31
11	S. dimorphus	31
12	Pediastrum braunii	32
13	P. muticum	33
14	P. tetras forma	33
15	P. angulosum	32
16	P. integrum	32
17	P. taylori	33
18,19	P. boryanum	32
20,24	P. tetras v. tetraodon	33
21	Tetradesmus sp. (cross-sectional view)	32
22	Scenedesmus bijuga	31
23	Pediastrum boryanum	32
25	Tetradesmus sp.	32

PLATE 3



PLATE 4

Figure		Page
1,3	Pediastrum boryanum v. undulatum	32
2	P. angulosum	32
4	Euglena spirogyra v. marchia	55
5	Coleochaete orbicularis	34
6	Lepocinclis fusiformis v. major	56
7	Microthamnion strictissimum	34
8,9	Protoderma viride	34
10	Trachelomonas horrida	56
11	Phacus sp.	56
12	Sorastrum spinulosum	33
13	Rhizoclonium crassipellitum	35
14	Lepocinclis fusiformis	56
15	Peridinium willei	57
16	Trachelomonas lacustris	56
17	T. bacillifera v. minima	56
18	Ulothris variabilis	34

PLATE 4



PLATE 5

Figure		Page
1	*Pleurotaenium trabecula* v. *maximum*	50
2,3	*P. ehrenbergii*	50
4	*P. trabecula* v. *elongatum*	50
5	*Gonatozygon brebissonii*	36
6,7	*G. aculeatum*	36
8,9	*Penium spirostriolatum*	49
10,15	*Ducellieria chodatii*	57
11	*Desmidium grevillii*	46
12	*Dinobryon cylindricum*	57
13,14	*Spirogyra gracilis*	35

PLATE 5



PLATE 6

Figure		Page
1	*Closterium libellula*	38
2	*C. striolatum*	38
3,4	*C. gracile*	37
5,6	*C. macilentum* v. *japonicum* forma	38
7,8	*C. rostratum*	38
9	*C. lunula* fa. *gracilis*	38
10	*C. intermedium*	37
11	*C. Dianae*?	37
12	*C. Dianae* v. *minor*?	37
13	*C. jenneri*	38
14	*Netrium digitus* v. *naegelii*	37
15	*N. digitus*	36
16	*N. interruptus*	37
17	*Tetmemorus laevis*	54

PLATE 6



PLATE 7

Figure		Page
1	*Cosmarium globosum*	41
2	*C. rectangulare* v. *hexagonium*	44
3,4	*C. pyramidatum*	43
5	*C. impressulum* ∝	41
6	*C. pseudoarctoum*	43
7	*C. impressulum* β	42
8	*C. laeve* v. *nov.* (?)	42
9	*C. subarctoum*	44
10	*C. abbreviatum* fa. *minor*	39
11	*C. pseudoquadratulum*	43
12	*C. hammeri*	41
13	*C. sexangulare*	44

PLATE 7



PLATE 8

Figure		Page
1,2	Cosmarium difficile v. dilatatum	41
3	C. venustum v. excavatum	46
4,5	C. pseudopyramidatum v. extensum	43
6	C. humile v. striatum	41
7,12	C. bioculatum	39
8	C. contractum v. ellipsoideum	40
9	C. quadratum fa. willei	43
10	C. pseudoprotuberans fa. minus	43
11	C. tenue v. depressum	45
13	C. subtumidum v. minutum	45
14	C. angulosum	39
15	C. undulatum v. minutum	45

PLATE 8



PLATE 9

Figure		Page
1	*Cosmarium ornatum* v. *perornatum*	42
2	*C. subcrenatum* v. *isthmochondrum* (See Plate 11, Figs. 1 & 2)	44
3	*C. botrytis* v. *tumidum*	39
4	*C. quinarium*	44
5	*C. portianum*	43
6	*C. reniform*	44
7-9	*C. ungerianum* fa.	45
10	*C. askenasyii* v. *americana*	39

PLATE 9



PLATE 10

Figure		Page
1	*Cosmarium margaritatum* fa. *subrotunda*	42
2	*C. amoenum*	39
3,4	*C. intermedium*	42
5	*C. crenatum*	40
6-11	*C. subspeciosum* v. *validius*	45

PLATE 10



PLATE 11

Figure		Page
1,2	Cosmarium subcrenatum v. isthmochondrum (See Plate 9, Fig. 2.)	44
3,4	C. sexnotatum v. tristriatum	44
5	C. novae-semliae v. granulatum	42
6	C. cymatonotophorum v. granulatum	40
7	C. trachypleurum v. fallax	45
8	C. bipunctatum	39
9,10	C. formulosum	41
11	C. connatum	40

PLATE 11



PLATE 12

Figure		Page
1	Euastrum oblongum	47
2	E. bidentatum v. speciosum	46
3	E. ansatum v. triporum	46
4	E. obesum	47
5	E. denticulatum v. angusticeps	46
6	Micrasterias rotata fa. evoluta	49
7	M. truncata v. neodamensis	49
8	M. americana	48

PLATE 12



PLATE 13

Figure		Page
1	Euastrum verrucosum v. perforatum	47
2	E. verrucosum v. rhomboideum	48
3	E. verrucosum v. vallesiacum	48
4	E. elegans	47
5	E. ansatum v. pyxidatum	46
6	E. didelta	47
7,8	E. gemmatum var. taftii	47

PLATE 13



PLATE 14

Figure		Page
1,8	Micrasterias denticulata v. angulosa	48
2,3	M. pinnatifida	49
4	M. rotata	49
5,6	M. conferta	48
7	M. muricata v. tumida	48

PLATE 14



PLATE 15

Figure		Page
1-3	*Staurastrum spongiosum*	54
4	*S. dickei* v. *circulare*	52
5	*S. subavicula*	54
6	*S. grande* v. *angulosum*	52
7	*S. pachyrhynchum*	53
8	*S. pyramidatum* fa. nov. (?)	54
9	*S. orbiculare*	53
10	*S. polonicum*	53
11	*S. punctulatum*	53
12	*S. polymorphum*	53
13	*S. brebissonii*	51
14	*S. margaritaceum*	53

PLATE 15



PLATE 16

Figure		Page
1,2	*Staurastrum anatinum*	52
3,5,6	*S. punctulatum* v. *kjellmani*	53
4	*S. acarides*	50
7,8,10	*S. sebaldii* v. *ornatum* (Figs. 7 & 8 are three-angled; Fig. 10 is four-angled)	54
9	*S. vestitum*	54

PLATE 16



PLATE 17

Figure		Page
1-3	Staurastrum breviaculeatum	51
4,5	S. kurilense fa. triquetra	52
6,7	S. kurilense	52
8	S. cuspidatum	52
9,11	S. alternans	51
10	S. crenulatum	51

PLATE 17



PLATE 18

Figure		Page
1	*Staurastrum arctiscon*	51
2	*Xanthidium subhastiferum*	55
3	*Staurastrum brevispinum*	51
4,5	*Cylindrocystis brebissonii*	36
6	*Spondylosium pulchellum*	50
7	*S. planum*	50

PLATE 18



PLATE 19

Figure		Page
1	*Aphanothece stagnina*	58
2	*Aphanocapsa elachista* v. *conferta*	58
3	*A. pulchra*	58
4	*Chroococcus prescottii*	58
5	*C. turgidus*	58
6	*Coelosphaerium pallidum*?	59
7	*C. kuetzingianum*	59
8,12	*Microcystis aeruginosa*	60
9	*Gloeothece rupestris*?	59
10	*Dactylococcopsis smithii*	59
11	*Merismopedia elegans*	60
13	*Eucapsis alpina* v. *minor*	59

PLATE 19



PLATE 20

Figure		Page
1	*Oscillatoria tenuis* v. *natans*	61
2	*O. agardhii*	61
3	*O. limnetica*	61
4	*Lyngbya aerugineo-caerulea*	60
5	*Oscillatoria tenuis*	61
6	*Phormidium corium*	61
7,8	*Tolypothrix distorta*	63
9	*Anabaena sphaerica*	62
10	*Cylindrospermum alatosporum*	62
11,12	*Calothrix epiphytica*	63
13	*Anabaena oscillarioides*	61
14-16	*Nostoc parmelioides* (Figs. 14 & 15 are peripheral filaments; Fig. 16 is an internal filament)	62
17	*Calothrix fusca*	64
18	*Nostoc parmelioides* (colony)	62
19	*Scytonema mirabile*	63
20	*Nostoc paludosum*	62

PLATE 20



GENUS AND SPECIES INDEX

Anabaena,
oscillarioides, 61; Pl. 20
sphaerica, 62; Pl. 20

Ankistrodesmus,
falcatus, 27; Pl. 1
gelifactum, 27; Pl. 1

Aphanocapsa,
elachista v. conferta, 58; Pl. 19
pulchra, 58; Pl. 19

Aphanothece,
stagnina, 58; Pl. 19

Asterococcus,
limneticus, 26; Pl. 1

Botrycoccus,
braunii, 29; Pl. 2

Bulbochaete, 35

Calothrix,
epiphytica, 63; Pl. 20
fusca, 64; Pl. 20

Chlamydomonas,
angulosa, 25; Pl. 1

Chroococcus,
prescottii, 58; Pl. 19
turgidus, 58; Pl. 19

Closterium,
dianae, 37; Pl. 6
dianae v. minor, 37; Pl. 6
gracile, 37; Pl. 6
intermidium, 37; Pl. 6
jenneri, 38; Pl. 6
libellula, 38; Pl. 6
lunula f. gracilis, 38; Pl. 6
macilentum v. japonicum, 38; Pl. 6
rostratum, 38; Pl. 6
striolatum, 38; Pl. 6

Coelastrum,
cambricum v. intermedium, 30; Pl. 3
microporum, 30; Pl. 3
printzii, 30; Pl. 3
proboscideum, 30; Pl. 3

Coelosphaerium,
kutzingianum, 59; Pl. 19
pallidum, 59; Pl. 19

Coleochaete,
orbicularis, 34; Pl. 19

Cosmarium,
abbreviatum f. minor, 39; Pl. 7
amoenum, 39; Pl. 10
angulosum, 39; Pl. 8
askenasyii v. americana, 39; Pl. 9
bioculatum, 39; Pl. 8
bipunctatum, 39; Pl. 11
botrytis v. tumidum, 39; Pl. 9
connatum, 40; Pl. 11
contractum v. ellipsoideum, 40; Pl. 6
crenatum, 40; Pl. 10
cymatonotophorum v. granulatum, 40; Pl. 11
difficile v. dilatatum f. nov., 41; Pl. 8
formulosum, 41; Pl. 11
globosum, 41; Pl. 7
hammeri, 41; Pl. 7
humile v. striatum, 41; Pl. 8
impressulum, 41; Pl. 7
intermedium, 42; Pl. 10
laeve, 42; Pl. 7
margaritatum f. subrotundata, 42; Pl. 10

novae-semliae v. granulatum, 42; Pl. 11
ornatum v. perornatum, 42; Pl. 9
portianum, 43; Pl. 9
pseudoarctoum, 43; Pl. 7
pseudoprotuberans f. minus, 43; Pl. 8
pseudopyramidatum v. extensum, 43; Pl. 8
pseudoquadratulum, 43; Pl. 7
pyramidatum, 43; Pl. 7
quadratum f. willei, 43; Pl. 8
quinarium, 44; Pl. 9
rectangulare v. hexagonium, 44; Pl. 7
reniform, 44; Pl. 9
sexangulare, 44; Pl. 7
sexnotatum v. tristriatum, 44; Pl. 11
subarctoum, 44; Pl. 7
subcrenatum v. isthomochondrum, 44; Pl. 9, 11
subspeciosum v. validius, 45; Pl. 10
subtumidum v. minutum, 45; Pl. 8
tenue v. depressum, 45; Pl. 8
trachypleurum v. fallax, 45; Pl. 11
undulatum v. minutum, 45; Pl. 8
ungerianum, 45; Pl. 9
venustum v. excavatum, 46; Pl. 8

Crucigenia,
rectangularis, 30; Pl. 3
quadrata, 30; Pl. 3

Cylindrocystis,
brebissonii, 36; Pl. 18

Cylindrospermum,
alatosporum, 62; Pl. 20

Dactylococcopsis,
smithii, 59; Pl. 19

Desmidium,
grevillii, 46; Pl. 5

Dictyosphaerium,
pulchellum, 29; Pl. 1

Dinobryon,
cylindricum, 57; Pl. 5

Ducellieria,
chodatii, 57; Pl. 5

Enallax,
alpina, 31; Pl. 3

Eremosphaera,
viridis, 27; Pl. 2

Euastrum,
anasatum v. pyxidatum, 46; Pl. 13
anasatum v. triporum, 46; Pl. 12
bidentatum v. speciosum, 46; Pl. 12
denticulatum v. angusticeps, 46; Pl. 12
didelta, 47; Pl. 13
elegans, 47; Pl. 13
gemmatum v. taftii, 47; Pl. 13
obesum, 47; Pl. 12
oblongum, 47; Pl. 12
verrucosum v. perforatum, 47; Pl. 13
verrucosum v. rhomboideum, 48; Pl. 13
verrucosum v. vallesiacum, 48; Pl. 13

Eucapsis,
alpina v. minor, 59; Pl. 19

Eudorina,
elegans, 25; Pl. 1

Euglena,
spirogyra v. marchica, 55; Pl. 4

Gloeocystis,
ampla, 26; Pl. 1
gigas, 26; Pl. 1
vesiculosa, 26; Pl. 1

Gloeothece,
rupestris, 59; Pl. 19

Golenkinia,
paucispina, 29; Pl. 2

Gonatozygon,
aculeatum, 36; Pl. 5
brebissonii, 36; Pl. 5

Kirchneriella,
lunaris v. irregularis, 27; Pl. 2

Lepocinclis,
fusiformis, 56; Pl. 4
fusiformis v. major, 56; Pl. 4

Lyngbya,
aerugineo-caerulea, 60; Pl. 20

Merismopedia,
elegans, 60; Pl. 19

Micrasterias,
americana, 48; Pl. 12
conferta, 48; Pl. 14
denticulata v. angulosa, 48; Pl. 14
muricata v. tumida, 48; Pl. 14
pinnatifida, 49; Pl. 14
rotata, 49; Pl. 14
rotata f. evoluta, 49; Pl. 12
truncata v. neodamensis, 49; Pl. 12

Microcystis,
aeruginosa, 60; Pl. 19

Microthamnion,
strictissimum, 34; Pl. 4

Mougeotia, 35

Netrium,
digitus, 36; Pl. 6
digitus v. nagelii, 37; Pl. 6
interruptum, 37; Pl. 6

Nitella, 55

Nostoc,
parmelloides, 62; Pl. 20
paludosum, 62; Pl. 20

Oedogonium, 35

Oocystis,
arctica, 28; Pl. 2
borgei, 28; Pl. 2
elliptica, 28; Pl. 2
parva, 28; Pl. 2
pusilla, 28; Pl. 2
sp., 28; Pl. 2

Oscillatoria,
agardhii, 61; Pl. 20
limnetica, 61; Pl. 20
tenuis, 61; Pl. 20
tenuis v. natans, 61; Pl. 20

Pandorina,
morum, 25; Pl. 1

Pediastrum,
angulosum, 32; Pl. 3 and 4
boryanum, 32; Pl. 3
boryanum v. undulatum, 32; Pl. 4
braunii, 32; Pl. 3
integrum, 32; Pl. 3
muticum, 33; Pl. 3
taylori, 33; Pl. 3
tetras, 33; Pl. 3
tetras v. tetraodon, 33; Pl. 3

Penium,
spirostriolatum, 49; Pl. 5

Peridinium,
willei, 57; Pl. 4

Phacus,
sp., 56; Pl. 4

Phormidum,
corium, 61; Pl. 20

Pleurotaenium,
ehrenbergi, 50; Pl. 5
trabecula v. elongatum, 50; Pl. 5
trabecula v. maximum, 50; Pl. 5

Protoderma,
viride, 34; Pl. 4

Quadrigula,
lacustris, 29; Pl. 2

Rhizoclonium,
crassipellitum, 35; Pl. 4

Scenedesmus,
bijuga, 31; Pl. 3
bijuga v. alternans, 31; Pl. 3

dimorphus, 31; Pl. 3
quadricauda, 31; Pl. 3

Scytonema,
mirable, 63; Pl. 20

Sorastrum,
spinulosum, 33; Pl. 4

Sphaerocystis,
schroeteri, 27; Pl. 1

Spondylosium,
planum, 50; Pl. 18
pulchellum, 50; Pl. 18

Spirogyra,
gracilis, 35; Pl. 5

Staurastrum,
acarides, 50; Pl. 16
alternans, 51; Pl. 17
anatinum, 52; Pl. 16
arctiscon, 51; Pl. 18
brebissonii, 51; Pl. 15
breviaculeatum, 51; Pl. 17
brevispinum, 51; Pl. 18
crenulatum, 51; Pl. 17
cuspidatum, 52; Pl. 17
dickiei v. circulare, 52; Pl. 15
grande v. angulosum, 52; Pl. 15
kurilense, 52; Pl 17
kurilense f. triquetra, 52; Pl. 17
margaritaceum, 53; Pl. 15
orbiculare, 53; Pl. 15
pachyrhynchum, 53; Pl. 15
polonicum, 53; Pl. 15
polymorphum, 53; Pl. 15
punctulatum, 53; Pl. 15
punctulatum v. kjellmani, 53; Pl. 16
pyramidatum, 54; Pl. 15
sebaldii v. ornatum, 54; Pl. 16
spongiosum, 54; Pl. 15
subavicula, 54; Pl. 15
vestitum, 54; Pl. 16

Tetmemorus,
laevis, 54; Pl. 6

Tetradesmus,
sp., 32; Pl. 3

Tetraëdron,
gracile, 26; Pl. 1

Tetraspora,
lamellosa, 25; Pl. 1

Tolypothrix,
distorta, 63; Pl. 20

Trachelomonas,
bacillifera v. minima, 56; Pl. 4
horrida, 56; Pl. 4
lacustris, 56; Pl. 4

Ulothrix,
variabilis, 34; Pl. 4

Xanthidium,
subhastiferum, 55; Pl. 18

Zoochlorella, 29

Zygnema, 36